



## Diseases of gerbera and its management

MB Navale<sup>1\*</sup>, RJ Sawant<sup>1</sup>, MS Wadikar<sup>2</sup>

<sup>1</sup> Department of Botany, Shri Muktanad College, Gangapur, Aurangabad, Maharashtra, India

<sup>2</sup> Department of Botany, Arts Commerce and Science College Kille Dharur, Maharashtra, India

### Abstract

Gerbera is commercially very important floriculture crop. There are multiple uses of gerbera hence this flower is in high demand at local and international market. Gerbera production is affected by various factors. One of the major devastating factors for gerbera production is different diseases occurring on gerbera. There is great need to manage these diseases effectively to increase production of gerbera all over the world. Effective management strategies ultimately help to reduce major losses in gerbera production.

**Keywords:** gerbera, diseases on gerbera, management of diseases

### Introduction

*Gerbera* is a genus of the family of Sunflowers with approximately 30 species and thousands of cultivar (Yeasmin and Shamsi 2013) [7]. The name gerbera was given in the honour of the German botanist and naturalist Traugott Gerber (1743) (Yeasmin and Shamsi 2013) [7]. The meaning of Gerbera is innocence and purity, this gerbera is also known by variety of names such as – African Daisy, Transvaal daisy and Barberton (Yeasmin and Shamsi 2013) [7]. Gerbera was first scientifically described as a South African species by J. D. Hooker in Curtis's Botanical Magazine in 1889 (Yeasmin and Shamsi 2013) [7]. Gerbera is very popular and commercially very important (Yeasmin and Shamsi 2013) [7]. Gerbera is one of the most commonly used cut flowers among the world (after rose, carnation, chrysanthemum, and tulip) (Yeasmin and Shamsi 2013) [7]. Long lasting, simple attractive structure and cheerful appearance make it suitable to combine with all other flowers (Yeasmin and Shamsi 2013) [7]. Gerbera is used as decorative garden plant, as cut flowers and also in floral arrangements in festivals, party and wedding ceremony (Yeasmin and Shamsi 2013) [7]. Now days, gerbera is cultivated as a garden plant throughout the India (Yeasmin and Shamsi 2013) [7].

### Diseases on gerbera

Gerbera plant is attacked by insects, pests, fungi, nematodes, bacteria, viral and phytoplasma pathogens (Yeasmin and Shamsi 2013; Meena *et al* 2015, Gautam *et al* 2020) [7, 3, 1]. Infections with these pathogens cause heavy losses in gerbera production (Meena *et al* 2015) [3].

### Fungal diseases on gerbera:

Fusarium crown rot (*Fusarium solani*) (Suneeta *et al* 2017), Fusarium root rot (*Fusarium oxysporum*) (Suneeta *et al* 2017), Phytophthora crown and foot rot (*Phytophthora cryptogea*) (Suneeta *et al* 2017), root rot (*Pythium irregulare*, *Rhizoctonia solani*) (Suneeta *et al* 2017), Pythium root rot (*Pythium spp.*) (Suneeta *et al* 2017), Rhizoctonia root (*Phytophthora cryptogea*), crown rot (*Phytophthora drechsleri*) (Suneeta *et al* 2017), Southern blight (*Sclerotium*

*rolfsii*) (Suneeta *et al* 2017), Collar rot (*Sclerotium rolfsii*) (Suneeta *et al* 2017) and leaf spot (*Alternaria alternata*) (Waghmare 2012) [6]. Crown and root rot (by *Phytophthora cryptogea*, *P. drechsleri*); Sclerotium rot (by *Sclerotium rolfsii*); Blight or Gray mould (*Botrytis cinerea*); Powdery mildews (by *Erysiphe cichoracearum*, *Oidium crysiphoides*); leaf spots (by *Alternaria alternata*) (Farhood and Hadian, 2012), (by *Corynespora cassiicola*) (Shi *et al.*, 2012) and by (*Phyllosticta gerberae*, *Alternaria spp.*). The downy mildews, yellow discoloration on leaf, later turning light to dark brown on gerberas are caused by *Bremia luctucae* (Wolcan *et al.*, 2010), and White rust (white erumpent sori) is caused by *Albugo tragopogonis* (Vazquez *et al.*, 1997).

### Bacterial diseases on gerbera

Bacterial leaf spot (*Pseudomonas cichorii*) is major bacterial disease on gerbera. Bacterial leaf spot disease on gerbera caused by *Pseudomonas cichorii* is also seen. The symptoms of this disease include: small to large spots, circular at first and then became irregular and dark brown to black spots (Ferreira, 1993) [9].

### Nematode diseases on gerbera

The major pests (whitefly, aphid, leaf miner, thrips, mites), diseases (powdery mildew, collar Rot, root rot, stem rot, leaf spot), nematodes (root-knot, spiral), and their symptoms, biology, spread, and management are known to create problems in gerbera (Reddy 2016) [10]. Different types of plant parasitic nematodes are found associated with gerbera elsewhere in the world (Lamberti *et al.*, 1987) [11], root knot nematodes belonging to *Meloidogyne spp.* are predominant in India (Nagesh and Parvath Reddy, 2001). In India, yield loss in gerbera due to *Meloidogyne incognita* was reported to be 31.1% (Nagesh and Parvatha Reddy, 2000) [12].

Different types of nematodes such as *Meloidogyne incognita*, *Helicotylenchus multicinctus*, *Pratylenchus coffeae*, *Tylenchorhynchus spp.* and *Rotylenchulus reniformis*.

### Management of Diseases

Application of integrated management system is more effective to control diseases on gerbera (Meena *et al* 2015) [3]. Leaf spot managed by Carbendazim and other fungicides (Waghmare 2012) [6]. It has been observed that Vermicompost incorporation at 20%, with or without chemical fertilizer, reduced the incidence of diseased plants, and the disease growth rate. Compared to all other treatments, the plant length, chlorophyll content, and number, length and diameter of inflorescences were also significantly higher with vermicompost 20% treatment, with or without chemical fertilizer (Rodriguez 2000). Soil solarization resulted in reduced root rot (root disease index 28.6%) in comparison to the untreated control (52.0%) 8 months after planting. Plants in the fumigated plots had 15.8% less disease than those in solarized plots (W. Kaewruang 1989)

### References

- Gautam KK, Kumar S, Raj SK. Diseases affecting gerbera cultivation and their control measures. An international journal of floriculture science and landscaping. The journal of the greens and gardens, 2020;3(5):1-16.
- Hausbeck MK. Disease management – managing powdery mildew on gerbera – The American floral endowment, 2003, 1-2.
- Meena KS, Ramya Bharathi SA, Raguchander T, Jonathan EI. *Meloidogyne incognita* and *Fusarium oxysporum* interaction in gerbera. African journal of Microbiology research (Academic journals), 2015;9(18):1281-1285.
- Salinas J, Glandorf DCM, Picavet FD, Verhoeff K. Effects of temperature, relative humidity and age of conidia on the incidence of spotting on gerbera flower caused by *Botrytis cinerea*. Neth J. Pl. Path, 1989;95: 51-54.
- Suneeta P, Eraivan AA, Nakkeeran S. Experiments on collar rot of gerbera by using biocontrol agents and fungicides in pot culture conditions. International journal of Agricultural science and research, 7(3), 481-486.
- Waghmare MB. Efficiency of mycotoxins of some plant extracts against *Alternaria alternata* (Fries) Keissler causing leaf spot of gerbera. Current Biotica, 2012;6(2):240-245.
- Yeasmin F, Shamsi S. Phylloplane Mycoflora of *Gerbera spp.* and their pathogenic potentiality, 2013;37(2):211-217.
- Bhat HA, Ahmad K.
- Ferreira-Pinto MM, Oliveira H. "A new bacterial disease of lettuce in Portugal caused by *Pseudomonas cichorii* (Swingle) Stapp." *Phytopathologia mediterranea*, 1993;32(3):249-253.
- Reddy PP. Gerbera. In: Sustainable Crop Protection under Protected Cultivation. Springer, Singapore, 2016. [https://doi.org/10.1007/978-981-287-952-3\\_29](https://doi.org/10.1007/978-981-287-952-3_29)
- Lamberti F, Tacconi R, Marinari A, Derrico FP, Basile M. "Major plant parasitic nematodes associated with flower crops in Italy and their control," *Difesa delta Pinate*, 1987;10:77-84.
- Nagesh M, Paravatha Reddy P. "Crop loss estimation in carnation and gerbera due to root-knot nematode *Meloidogyne incognita* (Kofoid and White) Chitwood," *Pest Management in Horticultural Ecosystems*, 2000;6:158-159.
- Rodríguez-Navarro JA *et al.* "The effect of vermicompost on plant nutrition, yield and incidence of root and crown rot of gerbera (*Gerbera jamesonii* H. Bolus)." *Fitopatologia*, 2000;35(1):66-79.
- Kaewruang W, Sivasithamparam K, Hardy GE. "Use of soil solarization to control root rots in gerberas (*Gerbera jamesonii*)." *Biology and fertility of Soils*, 1989;8(1):38-47.