



## *Nyctanthes arbor-tristis*: A traditional herbal plant with miraculous potential in medicine

Himanshi Rawat<sup>1</sup>, Yashika Verma<sup>1</sup>, Ayesha<sup>2</sup>, Neha Saini<sup>1</sup>, Neha Negi<sup>1</sup>, Hem Chandra Pant<sup>1</sup>, Aditi Mishra<sup>1</sup>, Mayank Singhal<sup>1</sup>, Arzoo Khan<sup>1</sup>, Naveen Gaurav<sup>1\*</sup>

<sup>1</sup> Department of Biotechnology, Shri Guru Ram Rai University, Dehradun, Uttarakhand, India

<sup>2</sup> Department of Biotechnology, Graphic Era Deemed To Be University, Dehradun, Uttarakhand, India

### Abstract

*Nyctanthes arbor-tristis* (NAT) Linn, also known as Parijata or Harsinghar is an Indian traditional plant of great importance. It has been used in Ayurveda, Siddha and Unani system of medicine for treatment of various infectious and non-infectious diseases. Each part of the plant has some medicinal value and is used for various pharmacological actions such as anti-arthritic, antispasmodic, antibacterial, anti-inflammatory, immunostimulant, antidiabetic, hepatoprotective, antioxidant, antimicrobial, anthelmintic, antileishmanial, anti-pyretic, anti-allergic, antiviral and CNS depressant. The phytochemical analysis of plant revealed presence of flavonoids, tannins, saponins, glycosides, alkaloids, steroids and phenolic compounds which are responsible for these pharmacological actions of the plant. The chemical drugs being used for treating various ailments are costly and moreover have various side effects. These drugs can be replaced by the herbal plants which have no such side effects and are cost efficient, and one such alternative can be NAT. This paper reviews recent researches, importance, applications, methods of propagation and future prospects of NAT. Particular emphasis has been given towards the applications and methods of propagation of the plant.

**Keywords:** *nyctanthes arbour*, *tristis*, Indian traditional plant, pharmacological actions, nat, propagation of the plant, etc

### Introduction

Various synthetic drugs are being used as medicine for treatment of diseases that comprise various side effects. Plants are source of various chemical substances that possess immense biological properties which can be used as alternative to these synthetic drugs.

Various plants were explored as source of bioactive agents that could treat different ailments. Among 4, 22, 000 flowering plants from the whole world, more than 50,000 plants are there which reportedly have medicinal and pharmacological use [1].

*Nyctanthes arbor-tristis* Linn. (NAT), a small divine ornamental tree, is used to pray God across India and is known for its fragrant white flowers. The plant is a well-known traditional Indian medicinal plant also, which in Ayurveda is used for various pharmacological actions such as anti-arthritic, antispasmodic, antibacterial, anti-inflammatory, immunostimulant, antidiabetic, hepatoprotective, antioxidant, antimicrobial, anthelmintic, antileishmanial, anti-pyretic, anti-allergic, antiviral and CNS depressant. It is an herbal remedy for treating sciatica, malaria, enlargement of spleen and other various infectious and non-infectious diseases. It is also used as blood purifier. NAT in local language is commonly known as Harsinghar, Parijata or night flowering jasmine [2, 3, 4]. The generic name 'Nyctanthes' came from the Greek words 'Nykhtha' meaning 'night' and 'anthos' meaning 'flower' while the specific name 'arbortristis' means 'the sad tree' because of the dull look of tree during the daytime [5].

The plant is native to the subtropical Himalayas of Nepal and India and geographically distributed in southern parts of India, northern Pakistan and in south-east Asian countries like Thailand, Malaysia and Indonesia [6]. In India, it is

distributed in the outer Himalayas and tract of Jammu and Kashmir, Nepal to East Assam, Bengal, Tripura and extended throughout the central region up to Godavari in south [1].

### Morphological Characters

It is a terrestrial woody shrub with white flowers having an extraordinarily strong and pleasant fragrance whole night that grows up to 10 meters in height and have a life span of 5-20 years [7]. The plant can be cultivated in tropical and subtropical regions all over the world, from sea-level up to 1500m altitude at the equator [6]. It usually grows in red and black soil with pH of 5.6-7.5 and prefers arid and semi-arid climatic conditions [1]. Flowering generally occur from July to October and flowers are arranged at the tip of branches or in axil of the leaves in cluster of 2-7 flowers together, each flower opening at dusk and falling at dawn. These flowers are fragrant and sessile having 6-8mm long, narrowly campanulate, hairy outside, glabrous inside, ciliated calyx and more than 13mm long, 6-8mm long tube, 5-8 unequally obcordate and cuneate lobed white corolla with an orange-red centre. Two stamens are inserted near the top of corolla tube and stigma is obscurely bifid. The fruits are flat, compressed, brown heart-shaped to round capsules with two sections each containing a single seed. Seeds are exalbuminous, testa are thick and outer layer of large transparent cells is heavily vascularized while cotyledons are flat, and radicle is inferior. The leaves are rough, hairy, decussately opposite, simple, 6-12 cm long, 2-6.5 cm wide with an entire margin. [8, 9].




Fig 1: Morphological Characters

**Taxonomical Classification**

**Kingdom:** Plantae  
**Subkingdom:** Viridiplantae  
**Infrakingdom:** Streptophyta  
**Superdivision:** Embryophyta  
**Division:** Tracheophyta  
**Subdivision:** Spermatophytina  
**Order:** Lamiales  
**Family:** Oleaceae  
**Genus:** Nyctanthes  
**Species:** Nyctanthes arbor-tristis

**Medicinal Usage of Various Parts of Nat In Ayurveda, Siddha And Unani System**

Nyctanthes arbor-tristis has been used by tribal people of India (Orissa and Bihar) along with its use in Ayurveda,

Siddha and Unani system of medicines for many years. Various pharmacological actions of different parts of plant have been investigated [10]. Seeds are used as anthelmintic, in alopecia, bilious pyrexia and powdered form is used for curing scurfy affections of scalp, piles and skin diseases [3]. Powdered stem bark is given to patients with rheumatic joint pain, used as an expectorant, in treatment of malaria, snake bite, bronchitis, ulcers, bleeding gums, anorexia, liver disorders, piles, worm infestation, blood disorder, oliguria, skin diseases and fever. Stem bark pounded with Zingiber officinale and Piper longum is boiled in water and the resultant liquid is taken for two days for treatment of malaria while the resultant paste on mixing with Arjuna bark is rubbed on the body for treatment of internal injury and joint broken bones. The oil prepared out of the bark of the plant is mixed with rice gruel and rock salt to prepare a

dose form (called anjana in ayurveda) used to apply in sclera of eye is found to be beneficial for various eye diseases [3, 6, 11]. Leaves are used for the treatment of various diseases such as obstinate sciatica, chronic pyrexia, rheumatism, intestinal worm infection, anorexia, haemorrhoid, liver disorders, biliary disorders, pyrexia with rigors, and as laxative, diaphoretic, diuretic and antidote to reptile venom. Leaf succulent mixed in honey, given thrice daily is effective in cough. Paste of leaves with honey is also given for treatment of pyrexia, high blood pressure, diabetes and leaf juice with honey is used to cure chronic fever and with loha-bhasma (mineral based ayurvedic medicine that contains elemental iron micro-fine particles prepared through the process of calcination under high temperature) is used in anaemia and hepato-biliary diseases (hepatic and spleen enlargement) while decoction of leaves is specifically recommended for obstinate sciatica. Bitter extract from leaves is given to children for expulsion of roundworms and threadworms and fresh leaves are used for preparing homeopathic medicines [3, 6, 8]. The juice of 3-7 leaves is used as blood purifier to prevent skin infection by tribal people in Orrisa [10].

Flowers are used as stomachic, carminative, astringent to bowel, antibilious, expectorant, hair tonic and treatment of haemorrhoid, obstinate remittent fever, sciatica, rheumatism, various skin diseases and for ophthalmic purposes [3, 6]. Decoction of flowers given orally to ward off wind in stomach, stimulate gastric secretion, improves expectoration of lungs and used in treatment of gout [12]. Roots are used for emaciation [4].

### Chemical Constituents of Nat




A variety of chemical constituent belonging to different classes such as terpenes, steroids, glycosides, flavonoids, alkaloids and aliphatic compounds were isolated and identified in various parts of NAT. Glycosides and






alkaloids, the secondary metabolites, are the largest group of chemicals produced by the plant. Glycoside and two alkaloids, one soluble in water and other in chloroform, was isolated from the bark. Roots are found to be composed of alkaloids, tannins and glucosides while leaves contain tannic acid, methyl salicylate, mannitol,  $\beta$ -sitosterol, flavanol glycosides, Astringent, Sugar, an amorphous glycoside 1%, mannitol 3%, amorphous resins 1.2%, ascorbic acid, oleanolic acid, nyctanthic acid, benzoic acid, Friedelin, lupeol, carotene and traces of volatile oil. It was found that frying the leaves in oil increases ascorbic acid content. Iridoid glycosides and arbortristosite A, B, D and E is found in seeds. Seed kernel yields 12-16% of pale yellow brown fixed oil that contains glucosides of linoleic, oleic, lignoceric, stearic, palmitic acid, 3,4-secotriterpene acid and  $\beta$ -sitosterol. Flowers consist of essential oil, nyctanthin, D-mannitol, tannin, glucose, carotenoids, glycosides,  $\beta$ -monogentiobioside ester of  $\alpha$ -crocetin (or crocetin-3),  $\beta$ -monogentiobioside,  $\beta$ -D monoglucoside ester of  $\alpha$ -crocetin,  $\beta$ -digiatiobioside ester of  $\alpha$ -crocetin (or crocin-1). Phytochemical analysis of ethanolic extract of flower revealed presence of renyolone, a cyclohexylethanoid, iridoid glycosides (6-o-trans-cinnamoyl-7-o-acetyl-6 $\beta$ -hydroxyloganin, arborside C, 6 $\beta$ -hydroxyloganin) and a phenylpropanoid glycoside, nyctanthoside. [8, 13, 14]. Flower oil consists of  $\alpha$ -pinen, p-cymene, 1-hexanol, methylheptanone, phenyl acetaldehyde, 1-decanol and anisaldehyde [15].






Bright orange corolla tubes of flower contain a colouring substance nyctanthin which is identical to  $\alpha$ -crocetin ( $C_{20}H_{24}O_4$ ) from saffron. The corolla tubes were formerly used for dyeing silk, sometimes together with safflower or turmeric [3].


### Some Phytoconstituents of Nat, Their Class and Structure

Table 1

S. No	Phytoconstituent	Chemical Class	Parts of plant constituting the phytochemical	Structure of the phytoconstituent
1.	Desrhamnosylverbacoside	Iridoid Glycoside	Leaf	
2.	$\beta$ -sitosterol	Steroid	Leaf, stem, seed	
3.	Friedeline	Terpenes	Leaf	

4.	Lupeol	Terpenes	Leaf	
5.	Nicotiflorin	Flavonoid	Leaf	
6.	Arbortristoside A, B, C, D and E	Iridoid glycoside	Seed	 <p style="text-align: center;">Arbortristoside A</p>  <p style="text-align: center;">Arbortristoside B</p>  <p style="text-align: center;">Arbortristoside C</p>

				 <p style="text-align: center;"><b>Arbotristoside D</b></p>  <p style="text-align: center;"><b>Arbotristoside E</b></p>
7.	6-o-trans-cinnamoyl-7-o-acetyl-6β-hydroxyloganin	Iridoid Glycoside	Flower	
8.	Arborside C	Iridoid Glycoside	Flower	
9.	6β-hydroxyloganin	Iridoid Glycoside	Flower	

10.	Oleanolic acid	Terpenes	Leaf	
-----	----------------	----------	------	---

### Pharmacological Applications of Nat

**Antioxidant activity:** Anowar *et al* in their work on the plant concluded the oxidative and protective role of hydro-alcoholic flower extract of NAT against the oxidative stress of hydrogen peroxide,  $H_2O_2$ .  $H_2O_2$  is weak oxidising agent that can penetrate the cell membrane and enter the cell where it reacts with  $Fe^{2+}$  and  $Cu^{2+}$  ions and forms hydroxyl radicals. These hydroxyl radicals further cause damage to the cell by interacting with the micro and macromolecules in the cell and inactivate enzymes usually by oxidation of the thiol (-SH) groups. They treated the lymphocytes isolated from chicken blood with  $H_2O_2$  that decreased the viability of cells by lowering the cellular antioxidant, reduced glutathione (GSH). The level of GSH was increased by 1.22-fold significantly when the lymphocytes were treated with the flower extract of NAT. At the same time, the specific activity of marker of membrane damage, lactate dehydrogenase (LDH) was also found to decrease as compared to untreated lymphocytes that suggests the non-toxic effect of extract on the cellular system. The experimental data of the study showed the antioxidant property of the crude extract of NAT [2]. Some other research has concluded leaves and stem of *Nyctanthes arbor-tristis* as a potential source of natural antioxidant. Phytochemical analysis of stem and leaves of the plant showed presence of flavonoids, tannins, saponins, glycosides, alkaloids, steroids and phenolic compounds. The antioxidant activity of NAT might be considered due to the phenolic compounds which act as free radical terminators [8].

**Antibacterial activity:** Antibacterial activity of flower, leaf, seed and fruit ethyl acetate and chloroform extract against gram-positive (*Staphylococcus aureus*) and gram-negative (*E. coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*) bacteria was examined by K. Priya *et al*, 2007. 300  $\mu$ l of both ethyl acetate and chloroform extract showed significant antibacterial activity against the bacteria tested. Flower ethyl acetate and seed chloroform extract showed broad spectrum antibacterial activity against gram-negative as well as gram-positive bacteria while leaf extract showed antibacterial activity against only but all the gram-negative bacteria. Fruit and seed ethyl acetate while flower and seed chloroform extract showed inhibitory effect against only *Pseudomonas aeruginosa* and *Klebsiella pneumoniae*. It was also found that the antibacterial activity of fresh plant parts was more than that of the dried one. Phytochemical analysis revealed presence of phytosterols, phenolics, tannins, flavonoids, glycosides and saponins. Phenolic compounds and tannin were found to be active against the bacteria. Tannins have been found to form irreversible complexes with proline rich proteins that results in inhibition of the cell protein synthesis and play an important role as stable and potent antioxidant, astringent and in treatment of diarrhoea and dysentery [4].

**Anti-fungal activity:** Different parts of NAT plant were examined for antifungal activity against three most prevalent clinical pathogenic fungi- *Aspergillus niger*, *Penicillium* and *Aspergillus flavus*. Fresh and mature leaves, seeds, stem, bark and flowers were collected and dried, and extraction was performed with distilled water, methanol and chloroform. The antifungal activity of the extracts was measured by well diffusion method in terms of “zone of inhibition” of fungal growth. The results revealed that only distilled water extract of stem and bark of NAT showed antifungal activity against *A. niger* only while chloroform extract of leaves was only effective against *A. flavus*. The study showed that the most effective results for antifungal activity was shown by methanolic extract of leaves, stem and bark of NAT against both *Aspergillus* and *Penicillium* [5].

**Antihistaminic and Anti-asthmatic activity:** Catalepsy is a condition where the limbs remain in whatever position they are placed. It is characterized by lack of response to the external stimuli and muscle rigidity and can be induced by neuroleptic drugs [16]. Experimental results proved that the clonidine-induced catalepsy can be inhibited by the petroleum ether extract of NAT bark. The effect of clonidine-induced catalepsy in Albino mice was studied and it was concluded that the cataleptic effect of clonidine in mouse is mediated by histamine from the mast cell and due to the mast cell stabilizing property of the NAT, this extract was able to inhibit the clonidine-induced catalepsy. The observation of this study indicated the antihistaminic activity of NAT bark and concluded that it may be useful in treatment of asthma [17]. Allergic asthma is a chronic inflammatory disease that affects breathing. It is characterized by bronchial obstruction reactions, airway inflammation and airway hyper-reactivity towards variety of stimuli like allergens, histamine, methacholine, etc. When exposed to allergen, the main component of the early asthmatic reaction is bronchoconstriction and it was concluded that under asthmatic conditions, the bronchial obstruction reaction and airway hyper-sensitivity are associated with the deficiency of nitric oxide production which plays a major role in maintenance of airway balance. The ethanolic extract of NAT leaves were examined for histamine-induced dose-dependent contraction of tracheal smooth muscles which concluded that the extract could inhibit the contractile responses produced by histamine and thus confirmed the bronchodilatory activity of the ethanolic extract of NAT leaves on airway smooth muscles. The results of experiment showed that the acting mechanism of the extract is through increase in production of nitrogen oxide, NO.

It has also been concluded that the anti-asthmatic and anti-allergic activity of NAT leaves is due to the presence of  $\beta$ -sitosterol [18].

**Wound healing activity:** The wound healing activity of NAT was tested on Wistar albino rats by Matadeen *et al.*, 2011. The rats were treated with 2% w/w NAT methanolic extract for 16 days. It was found that the complete epithelization of both incision and excision wounds took about 16 days and wounds were completely healed. It was concluded that 300mg dose/kg of NAT extract can be considered a good solution for healing both the types of wound [19].

**Immuno-stimulant activity:** Aqueous leaf extract of NAT has been found to be a potent immunomodulator which is evidenced by both humoral and cell-mediated responses. Flower has also shown immune-stimulant activity that activates the cell-mediated immune system. Immunomodulator activity against systematic candidiasis in mice was shown by the seed and root ethanolic extracts of NAT. Both iridoid glucosides i.e., arbortriosides A and C were isolated from seeds of the plant [7, 8].

**Anti-ulcerogenic and ulcer-healing property:** Peptic ulcer, one of the major gastrointestinal disorders, occur due to imbalance between offensive (mainly acid, pepsin, *H. pylori* and bile salts) and defensive (mainly involves mucus-bicarbonate secretion and prostaglandin) factors. Reduction in production of gastric acid and gastric mucosal protection are the two major approaches for gastric ulcer disorder therapy. Arbortrioside-A and 7-O-trans-cinnamoyl-6 $\beta$ -hydroxyloganin from NAT have been reported to possess anti-ulcerogenic and ulcer-healing property. These two not only prevent the formation of irritant-induced gastric ulcers but also increase gastric ulcer healing [20].

**Analgesic and anti-inflammatory activity:** The pathophysiological response of mammalian tissue towards various hostile agents such as infectious organisms, toxic chemicals, physical injury or tumor growth is called inflammation. Edema formation, leukocytes infiltration and granuloma formation are components of inflammation. Glucocorticosteroids and nonsteroidal anti-inflammatory drugs (NSAIDs) are the two main types of anti-inflammatory agents but due to their adverse side effects various alternatives of NSAIDs and opiates are being searched all over the world. NAT was examined for the same and experiments were carried out on various animal models. The methanolic extract of NAT stem bark showed the analgesic and anti-inflammatory activity by preventing stimulation of nociceptive components which may happened due the inhibition of the production of prostaglandin and related compounds. Experimental results also showed that the extract reduced the rate of edema in carrageenan-induced rat paw edema model [21].

The peripheral analgesic activity of leaf extract of NAT was studied in mice using the acetic acid induced writhing test. Studies revealed that the presence of flavonoids in NAT might be responsible for the analgesic activity of the plant. Flavonoids are known for inhibiting synthesis of prostaglandin which are involved in pain perception by inhibiting cyclooxygenase enzyme. Other chemicals like methyl salicylate and benzoic acid which have strong analgesic effect are also found in NAT [22].

**Anthelmintic activity:** Parasitic diseases, also known as parasitosis, are infectious diseases caused or transmitted by

parasites. Helminths of class Cestoidea and Trematoda are endoparasites (live inside body of the host) and cause helminthiasis, also called worm infection. They often live in the gastrointestinal tract of their hosts but may also burrow into the organs inducing physiological damage.

Anthelmintics are the agents which are used to expel out the parasitic worms (helminths) from body, either by stunning or killing them. Alcoholic and aqueous extract of NAT bark was examined for their anthelmintic activity against *Pheretima posthuma*. *P. posthuma* due to its anatomical and physiological resemblance with the intestinal worm of human and easy availability was selected for this assay. The study revealed that the alcoholic extract was not only able to paralyze the worm in 8.53 min but also caused death in 13.05 min. Phytochemical analysis revealed presence of tannin that was responsible for the anthelmintic activity of NAT. Tannin can bind to the free protein present in gastrointestinal tract of host animal or glycoprotein present on cuticle of the parasite and may cause death of the worm [11].

**Antileishmanial activity:** In a study, iridoid glucosides *viz* arbortrioside A, B, C and 6 $\beta$ -hydroxy-loganin isolated from NAT were reported for the antileishmanial activity. In another study, calceolarioside A isolated from the methanolic extract of NAT leaves by bioactivity-guided fractionation showed in-vitro antileishmanial activity, which was determined to be IC<sub>50</sub> = 20 $\mu$ g/mL while its in-vivo efficacy, when it reduced the hepatic and splenic parasite burden by 79% and 80% respectively, was noted to be 20mg/kg body weight in *L. donovani* Ag83 infected golden hamster model. This was the first report isolating calceolarioside A from NAT and describing potent activity of NAT against visceral leishmaniasis [3].

**Anti-viral activity:** Anti-viral activity of NAT was examined against encephalomyocarditis (EMCV) and Semliki Forest Viruses (SFV) in Swiss albino mice. The experimental results showed that the crude ethanolic extract, n-butanol fraction and isolated iridoid glycosides (Arbortriosides A and C) showed inhibition of 75% of cytopathic effect caused by both the viruses. In-vivo studies against EMCV at different doses showed that the crude ethanolic extract and n-butanol fraction protected 40% of animals infected with EMCV at 250mg/kg body weight dose while the aqueous fraction protected 50% of animals infected with EMCV at 125 mg/kg body weight dose. Whereas in case of in-vivo anti-SFV activity the result was most promising with n-butanol fraction that provided 60% protection to animals infected with SFV at 125 and 62.5 mg/kg body weight dose and Arbortrioside A that also protected 60% of SFV infected animals but at a much lower dose of 31.2mg/kg body weight. Both these fractions and isolated compounds did not produce any significant antiviral activity when administered orally. Further study revealed that with increasing dose of virus (SFV) the level of protection declined. However, animal treated with Arbortrioside A recorded 66, 65 and 50% protection against 5, 10 and 100 LD<sub>50</sub> concentration. Based on the experimental results, it was concluded that the arbortrioside A isolated from n-butanol fraction of NAT possess maximum antiviral activity against enveloped virus (SFV) while moderate antiviral activity of n-butanol fraction was observed against EMCV. It was also observed that the

antiviral activity of n-butanol and arbortristiside A and C depends upon the route of administration, challenged virus concentration and dose of the compound [23].

**Hypoglycemic and hypolipidemic activity:** Diabetes mellitus is a major global disease from which millions of people are suffering. In diabetes, it is important not only to control blood glucose level but also the blood lipid level as diabetes tends to increase low-density lipoprotein cholesterol and decrease high-density lipoprotein cholesterol levels in blood that triggers coronary occlusions and blocks. It has been concluded that the present treatment of diabetes using synthetic hypoglycaemic agents may cause adverse effects resulting in hypoglycaemia, gastro-intestinal disturbances, renal toxicity and hepatotoxicity and this is why plants are been considered as an alternative. Different doses of boiled aqueous extract of fresh NAT flowers were examined for the hypoglycaemic and hypolipidemic activity in mice. 200, 500 and 750mg/kg of the extract was administered in mice and subsequently fasting and random glucose concentration were examined. After oral administration of the extract, the effect of extract on glucose tolerance, gastrointestinal glucose absorption, liver and skeletal muscle glycogen content, diaphragm uptake, serum lipid profile and in-vitro amylase assay were examined in mice. The experimental results concluded that 500 and 750 mg/kg of the extract reduced fasting glucose levels by 49% and 39% respectively at 4hr post-treatment while 500 mg/kg of the extract decreased the random blood glucose level significantly by 32% at 4hr post treatment. The extract also significantly inhibited glucose absorption from intestine by 85%, increased diaphragm uptake of glucose by 64%, decreased level of total cholesterol and triglycerides by 44.8% and 53% respectively, and increased the high-density lipoprotein cholesterol by 57%. It also exhibits inhibition of  $\alpha$ -amylase enzyme activity by 16.66%. The biochemical and toxicological effect of the extract neither showed any deaths nor there were any signs of clinical toxicity, stress and aversive behaviour during treatment period. Further histopathology analysis of liver and kidney portion after treatment with extract did not show any effect. The experimental data concluded that the boiled aqueous extract of fresh NAT flowers possess hypoglycaemic and hypolipidemic activity which can be used as an alternative medicine for treatment of diabetes and toxicological results considered it safe for oral administration [24].

**Antihyperlipidemic activity:** Hyperlipidaemia is increased level of lipid in blood which is first and for most factor responsible for diseases like atherosclerosis, coronary heart disease, ischemic cerebro-vascular disease, hypertension, obesity and diabetes mellitus (Type II) etc. Methanolic extract of NAT leaves was examined for antihyperlipidemic activity in Wistar albino rats. The extract at the doses of 200 and 400 mg/kg body weight showed significant decrease in lipid profile like triglycerides, total cholesterol (TC), low density lipoprotein (LDL), very low-density lipoprotein (VLDL), and significant increase in high density lipoprotein (HDL). The effect of lipid lowering might be result of plant sterols ( $\beta$ -stigmasterol and  $\beta$ -sitosterol) which reduce cholesterol absorption and thus increase faecal excretion of steroids and thus results in decrease of body lipids. It was concluded that the methanolic extract of NAT leaves showed antihyperlipidemic activity with specific and non-

specific mechanism which might be due to the presence of phytochemicals like phenol, triterpenoids and flavonoids in the extract [25].

**Anti-arthritis activity:** Arthritis is an autoimmune condition of synovial joints, triggered generally by inflammatory mediators and infections which is characterized by chronic inflammation that results in pain, inflammation of synovial joints, pannus formation, cartilage rupture, impaired movement and disability. Currently for treatment of arthritis nonbiologic disease modifying antirheumatic drugs (DMARD) (such as methotrexate), nonsteroidal anti-inflammatory drugs (NSAID) (such as piroxicam), biological therapies (such as inhibitors of IL-6, IL-1 and TNF- $\alpha$ ) and glucocorticoids (such as methyl prednisolone and triamcinolone) are used. NSAID are considered as the first line therapy of rheumatoid arthritis that effectively attenuate swelling, pain and joint stiffness but have adverse side-effects including stomach ulcers, bleeding, dyspepsia and risk of cardiovascular problems. Non-biologic DMARDs used for treatment can cause reversible alopecia, rash, nausea, loss of appetite, elevated formation of rheumatoid nodules, mouth ulcers and neurological problems while anti-TNFs therapy are responsible for abdominal pain, headache, itching, bruising, bleeding, rash, vomiting, diarrhoea, injection site reaction, cellulitis and respiratory tract infections. Chronic use of glucocorticoids also has multiple adverse effects including increased risk of osteoporosis, diabetes mellitus, peptic ulcer, gastrointestinal bleeding, cataract and infection. All the adverse effects related to the current treating therapies, increasing percentage of rheumatoid arthritis patients are going towards the alternative medicines. Methyl, ethyl acetate and n-hexane extracts of mature leaves of NAT were examined for anti-arthritis activity using rat model of FCA-induced arthritis. The comparative study of extracts showed that the ethyl acetate extract of NAT leaves was most promising than others. The ethyl acetate extract showed maximum inhibition of paw edema as compared to the other two extract. It more significantly attenuated infiltration of inflammatory cells in ankle joint, total leukocyte count and reduction in bone erosion as compared to other two as well as the reference drug, piroxicam. The phytochemical analysis of ethyl acetate extract showed presence of phytol, eugenol and  $\alpha$ -terpineol. Phytol suppresses oxidative stress and cytokine production playing anti-inflammatory role while eugenol possess anti-inflammatory activity via regulating redox reaction.  $\alpha$ -Terpineol is also known for its anti-inflammatory activity that inhibits proinflammatory IL-6 receptor gene expression levels whereas terpenoids regulate inflammatory and immune responses by inhibiting nuclear factor kappa B. The antiarthritis activity of the extract was attributed to the presence of these terpenes and terpenoids. The hepatotoxic and nephrotoxic effects of NAT extracts were evaluated that showed absence of any such effects in group of animals treated with the extracts and considered safe [26].

**Anti-spasmodic activity:** A study taken on NAT for antispasmodic activity of the plant was estimated using guinea pig ileum preparation against acetylcholine, a spasmodic agent. Antispasmodic activity of ethanolic extract of different parts of plant was estimated on the basis of inhibition of contractile effect of acetyl choline by

various dilution of the extract, out of which flower and seed extract were the most effective. The contractile response of 0.0002mg of acetylcholine was inhibited by 72mg of flower and 90mg of seed extract which was less than that of 16mg piperazine citrate [27].

**CNS depressant action:** A study was carried out with water soluble portion of ethanol extracts of flowers, barks, seeds and leaves of NAT to evaluate CNS depressant activity of plant. It has been already concluded that the leaves of the plant possess hypnotic and tranquilizing activity whereas its flower possess sedative activity. The ethanol extract of plant was extracted by Soxhlet extraction and the extract was examined for the pharmacological activity in adult male Swiss mice. The CNS depressant activity was evaluated by observing the prolongation of sleeping in mice induced by pentobarbital sodium. The attempts were made to determine the possible mechanism behind the activity by determining their effect on brain monoamine neurotransmitters like serotonin and dopamine. The study showed that the ethanolic extract of leaves, flowers, seeds and bark (600 mg/kg) of NAT possess significant CNS depressant activity and some muscle relaxant activity. The leaves, flowers, seeds and bark showed significant and dose dependent prolongation of the onset and duration of sleep, which was comparable to chlorpromazine, the standard drug. The highest CNS depressant activity was observed to be possessed by the leaves. It was also concluded that the activity possess by the extract might be due to the decrease in dopamine and increase in serotonin level in brain [28].

**Hepatoprotective activity:** Petroleum ether and methanol extract of NAT bark were examined for the hepatoprotective activity using carbon tetrachloride (CCl<sub>4</sub>) induced hepatotoxicity in Swiss albino mice. The hepatic toxicity was induced in the liver of mice by injecting CCl<sub>4</sub> and the level of Serum Glutamate Pyruvic Transaminase (SGPT), Serum Glutamate Oxaloacetic Transaminase (SGOT), Alkaline phosphate (ALP), Direct Bilirubin (DB) and Total Bilirubin (TB) in the serum of mice was measured for determining the function of liver. Silymarin was used as standard drug for the study. The administration of Silymarin and both the extracts (100 mg/kg and 200 mg/kg) significantly decreased the serum marker enzymes (SGPT, SGOT, ALP, DB & TB). The dose of 200 mg/kg of methanol extract was found to be more effective in comparison to that of 100mg/kg methanol and 100 mg/kg and 200 mg/kg of petroleum ether extract of NAT. The phytochemical analysis of petroleum ether and methanol extract of NAT showed presence of alkaloid, cardiac glycosides, tannin, saponin, terpenoid, phlobatannins, fixed oils, fats and flavonoids that might be responsible for the activity [29].

**Anti-pyretic activity:** Fever is a complex physiological response which is triggered by infectious or aseptic stimuli that results in elevated body temperature which occur due to increase in concentration of prostaglandin E (2) in certain parts of brain that further alter the firing rate of neurons that control thermoregulation in hypothalamus [30]. Antipyretics are drugs that reduce the elevated body temperature. NAT was examined for antipyretic activity in albino rats for potential on normal body temperature and yeast-induced pyrexia. The whole plant extract of NAT showed significant

antipyretic activity at the dose of 200 mg/kg by reducing normal body temperature and yeast-provoked elevated temperature in dose dependent manner. It was also found that the effect of whole plant extract of NAT dose was comparable to that of paracetamol (150 mg/kg), a standard antipyretic agent [31]. In another study the petroleum ether and methanol extracts of NAT bark were examined for the antipyretic activity of the plant against yeast induced pyrexia in mice. Fever was induced by 20% aqueous solution (10 mg/kg) of Brewer's yeast subcutaneously injected into the dorsum region of mice and aspirin (150 mg/kg) was used as standard drug. It was found that both petroleum ether and methanol extract (100 mg/kg and 200 mg/kg) showed antipyretic activity by significantly reducing rectal temperature after its administration [29].

**Anxiolytic activity:** Hydroalcoholic extract of NAT leaves was examined in adult Albino rats and Wister mice for its anxiolytic activity using various models like elevated zero maze, elevated plus maze, open field exploratory behaviour, novelty induced suppressed feeding test and social interaction test. Extract was given orally to rats in two different doses i.e., 250 and 500 mg/kg that showed dose dependent significant increase in open field ambulation, rearing, self-grooming and activity in centre manifesting anxiolytic activity of NAT. However, there were no significant changes in open field fecal droppings. In open field, animals are in novel environment, due to anxiety and fear, they express increased defecation and decreased ambulation, exploration, freezing, rearing and grooming behaviour and these behavioural changes are attenuated by classical anxiolytics and augmented by anxiogenic agents. Similarly, anxiolytic agents increase responses in elevated plus maze and elevated zero maze tests, open closed arm entries and time ratios provide a measure fear induced inhibition of exploratory activities. In experimental results, it was found that NAT treated rats exhibited dose dependent significant increase in time spent in open arms and entries made on open arms whereas there was significant decrease in time spent in enclosed arms and entries on enclosed arms as compared to the control. The extract also increased social interaction and attenuated novelty induced feeding latency in comparison to vehicle treatment. Lorazepam (500 mg/kg; i.p.) was used as standard drugs for comparison which showed significant anxiolytic activity and the effects were found to be more than that of the NAT extract. Overall experimental results provided positive result for the anxiolytic activity of the hydroalcoholic extract of NAT leaves [32].

**Anti-amnesic activity:** Neurological disorders are more common disorder in population aging between 18 and 60 years, characterized by various psychiatric disorders like depression, panic attacks, phobias, generalized anxiety and obsessive-compulsive and post-traumatic disorders. Among these, Alzheimer's disease is the most common degenerative disease which is characterized by neural loss, inflammation, memory loss and age-related decline in cognitive and learning disabilities. The pathogenesis of Alzheimer's disease results due to dysfunction of cholinergic system and increased oxidative stress in the hippocampal part of the brain. The anti-amnesic activity of ethanolic flower extract of NAT was studied and the anti-memory impairment activity of the extract was confirmed

using Elevated plus maze (EPM), Passive avoidance (PA) and Morris water maze (MWM) tests in adult male Wistar rats. The study concluded that the ethanolic extract of NAT flower improves memory by inhibiting acetylcholinesterase activity and reducing the oxidative stress in the brain tissue. GC-MS and UPLC-Q-TOF-MS/MS were used for identification of the biologically active compound present in the extract which regulate the brain function. Phytol and Loliolide were identified through GC-MS analysis while four compounds, namely 4-coumaric acid, chlorogenic acid hemihydrate, chalcone and melatonin were identified using UPLC-Q-TOF-MS/MS analysis in the extract. These active compounds were reported first time in the flower of NAT and neuroprotective activity of all these compounds was reported in different study, concluding that the extract might be useful in treating and controlling several neurodegenerative diseases [33]. In another study, the hydroalcoholic extract of NAT leaves was investigated for its effect on scopolamine induced amnesia using Morris water maze test in Albino Swiss male rats. After injecting 0.4 mg/kg scopolamine rats showed impairment of memory compared to the control group in which little change in escape latency time was noted to find hidden platform. Plant extract was administered in different doses and the decrease in escape latency time to find the hidden platform and the time spent by rats in the target quadrant was taken as the index of retrieval. Transfer latency time was estimated using elevated plus maze and it was found that the extract increased the transfer latency time, concluding that plant have a nootropic activity. Further the in-vivo study of NAT supported by the biochemical parameter estimation such as brain Acetyl cholinesterase activity, Glutathione estimation, malondialdehyde estimation and nitric oxide estimation concluded that the hydroalcoholic extract of NAT leaves improve learning and memory [15].

**Anti-cancer activity:** The methanolic crude extract of leaf, fruit and stem of NAT were examined in-vitro for antioxidant and anticancer activity of plant. By DPPH free radical scavenging assay antioxidant activity of extract was evaluated, it was reported that the dried fruit methanol extract of NAT showed 93.8% scavenging effect of phenolic crude at 1000 mg/ml conc., dried stem methanolic extract showed a moderate value of 69.9% at 100 mg/ml conc. while least value was 27.8% that was observed with dried leaves methanol extract at 1.0 mg/ml conc. Based on antioxidant activities the anti-cancer activity was evaluated by MTT assay on MDB MB-231 cancer cell lines. It was found that out of all the extracts, dried fruit methanol extract showed high degree of inhibition against human breast cancer cell lines (MDA-MB 231). The phytochemicals isolated from NAT dried fruit methanol extract were glycosides, tannins, phenols and steroids which were predicted to be responsible for the anticancer activity of the plant [34]. In another study, the methanolic extract of leaves of NAT were examined for anti-cancer activity of the plant against Ehrlich Ascites Carcinoma (EAC) cells ( $10^7$  cells/mouse) where the extract was injected into intraperitoneal route (200 and 400 mg/kg body weight) of Swiss Albino mice. For the study 5-Fluorouracil (20mg/kg body weight) was injected through intraperitoneal route as standard anti-cancer drug. Based on the ability to inhibit cancer cell growth in ascitic fluid of mice, the anticancer activity of the extract was evaluated. Various parameters

were taken to establish the potency of the anticancer property of methanol extract of NAT leaves including percentage inhibition of total cell count, tumour volume, viable and non-viable cell counts; percentage increase in life span and hematological parameters. Administration of extract at the doses of 200 and 400mg/kg b.w. significantly reduced the total cell count and tumour volume. At 200 mg/kg conc. the percentage of inhibition of total cell count and percentage inhibition of tumour volume was observed to be 84.75% and 75.52% respectively while at 400 mg/kg conc. the percentage of inhibition of total cell count and percentage inhibition of tumour volume was observed to be 90.36% and 96.37% respectively. Administration of extract at doses of 200 and 400 mg/kg also significantly decreased the viable cell count and level of WBC while increased non-viable cell count, life span and the level of both RBC and haemoglobin. The overall results of the study concluded that the methanol extract of NAT leaves possess anticancer activity [35].

### Method of Propagation

Due to its peculiar and pleasant fragrance it is often cultivated in garden. The shrubs can be propagated by seeds as well as by cutting, but the phenolic compounds leaching out of the imbibed seed decreases the rate of germination by seeds [12]. Due to low seed setting, poor germination by seed because of the inhibitory compounds and attack of insects during ripening stage, many young seedlings die under natural conditions and thus propagation of NAT by seed becomes very difficult [37, 38]. The inhibitory phenolic compounds are stored in the pericarp of the seed coat. It was studied that by either removing both the coverings or treating seeds with a solution of antioxidants like polyvinylpyrrolidone (PVP) and polyvinylpolypyrrolidone (PVPP) prior to germination can improve germination rate [12]. In the Red Data Book, NAT has been recorded as a 'valuable plant' which is critically endangered and at extremely high risk of extinction in the wild. Therefore, it is highlighted for conservation in many documentations for which in-vitro propagation is been used. The natural propagation of NAT is also a slow process that could not fulfil market demand [36]. The plant is being markedly diminishing due to destruction of its natural environment, excessive over exploitation and unresolved problem of seed viability and poor germination [39]. Hence, for commercial, ornamental and conservation purposes studies are needed to be conducted on production of the plant in large scale [38]. For saving this species from extinction, it is required to propagate these plants using alternative approaches like tissue culture techniques [39]. Plantlets of NAT can be raised by in vitro methods using excised cotyledons, hypocotyls, roots, leaves and bases of internodes that are able readily form callus in the culture. However, calli from cotyledons, hypocotyls and roots exhibited faster growth as compared to those from leaves and internodal bases when cultured in Murashige and Skoog's (MS) media with 2, 4 dichlorophenoxyacetic acid (2,4D), Naphthalene acetic acid (NAA) and coconut milk [12].

### In-vitro method of propagation of NAT

In a study conducted by Mahipal *et al* on in-vitro regeneration of NAT, the shoot tips and nodal segments with axillary buds were used as explant for establishment of culture. The explants were passed through surface

sterilization protocol where they were thoroughly rinsed under running tap water, then were pre-soaked in Tween-20 (liquid detergent) for about 15min., and then were treated with 0.1%(w/v) Bavistin (a fungicide) for 5min. After that, the explants were soaked in 0.1% (w/v) HgCl<sub>2</sub> for 5-6min in Laminar Air Flow cabinet, where they were finally rinsed with sterile distilled water for 8-10 times. Murashige and Skoog's basal medium was used for shoot regeneration and proliferation which was supplemented with 30g/L sucrose, 8g/L agar and additives (50mg/L ascorbic acid, 25mg/L each of adenine sulfate, citric acid and arginine). Before sterilizing the medium in the autoclave at 121°C for 15 min the pH of the medium was adjusted to 5.8 ± 0.02 using 0.1 N NaOH or HCl. For shoot propagation, the medium was augmented with cytokinin, 6-benzylaminopurine (BAP) or kinetin alone or in combination with Indole-3 acetic acid (IAA) (0.1-0.2 mg/L BAP or kinetin and 0.5 mg/L IAA). The proliferated shoot with mother explants were transferred to fresh medium of same conc. of growth hormone for 2-4 passages to check significance of subsequent transfer on shoot multiplication. The cultures were incubated for a photoperiod of 12/12hr light/dark. 5-6 cm long and healthy in-vitro regenerated shoots were then excised and implanted to full-, half- and one-fourth MS salts medium augmented with different conc. (1.0-5.0 mg/L) of auxin (Indole acetic acid, IAA or indole-3-butyric acid, IBA or 1-naphthaleneacetic acid, NAA) at 50, 100 and 200 mg/L activated charcoal for induction of root. The cultures were further incubated with diffused light for photoperiod of 8/16 hr light/dark. After 4-5 weeks, the in-vitro rooted microshoots were then taken out from the culture vessels and after washing thoroughly to remove adhered gel on roots the microshoots were transplanted to paper cups containing 55g autoclaved Soilrite (mixture of perlite, Irish peat moss and exfoliated vermiculite) moistened with 1/4<sup>th</sup> MS salts solution. The entire setup was maintained in greenhouse with 80% relative humidity and plants were watered at an interval of days with 1/4<sup>th</sup> MS inorganic solution. The following results were evaluated after the study:

1. After 10 days of culturing shoot tips and nodal segment of NAT on MS medium with 3.0 mg/L BAP, three to four microshoots were produced and the number shoots was increased to nine when the explants were incubated for 4weeks on MS medium with same conc. of growth regulators.
2. Kinetin alone and in combination with BAP (BAP + kinetin) were not much effective in induction of shoot as compared to the results of BAP alone.
3. Bud break and growth of axillary meristem cannot be obtained with media devoid of growth regulators.
4. Nodal shoot with axillary buds gave better results in comparison to shoot tip as explant.

5. The medium containing cytokinin with IAA (0.5mg/L) produced multiple shoots in comparison to the medium containing cytokinin only.
6. The rooting of in-vitro shoots was achieved in the presence of auxin on half strength MS medium with 50mg/L charcoal and maximum root regeneration was obtained with NAA (3.0mg/L) within 4 weeks while the lower number of roots per shoots were observed with the same concentration of IAA and IBA.
7. The explant collected during November to January showed great regenerative effect on shoot induction showing that season of collection of explant plays an important role in success of micropropagation.
8. Within 6 weeks, the in-vitro rooted plantlets got adapted to the natural environment of greenhouse and a survival rate of about 86% was recorded during the hardening process <sup>[36]</sup>.

In another study conducted to develop a simple and high frequency regeneration protocol for NAT, the influence of Putrescine was evaluated as a regeneration enhancer in combination with cytokinin and auxin. It was observed that putrescine at a conc. of 50mg/L had maximum promoting influence on percentage of culture response (93.33%) and number of shoots regenerated per explant, but both the percentage of culture response and number of shoots regenerated per explant decreased when conc. of putrescine was increased after 50mg/L. It was assumed that when the endogenous pools of polyamines (e.g., putrescine) are at minimum level they may lead to formation of methionine that in turn produces ethylene and when putrescine is at optimum level (50mg/L) organogenic response might be induced by them.

Whereas when putrescine is at higher level, they may divert to other biosynthetic pathways like ethylene or may cause feedback inhibition. Well elongated microshoots obtained when were subjected to in-vitro rooting, 86.66% rooting was observed and after hardening and acclimatization for 6 weeks, plants were successfully rooted in natural environment and a survival rate of almost 92% was obtained. The results showed that the regeneration frequency of NAT was enhanced by putrescine by efficiently modulating ethylene and therefore putrescine can be used as an additive in culture media to increase shoot regeneration <sup>[39]</sup>.

#### **Generalized steps for in-vitro propagation of NAT by nodal shoot**




Fig 2

### Conclusion and Future Prospect

Various synthetic chemicals are being widely used for treatment of different diseases but encompass adverse side effects. Due to these side effects of the drugs, various alternatives are being explored by the researchers and for the same plants are being studied. One such plant is *Nyctanthes arbor-tristis* (NAT). The broad-spectrum medicinal use of NAT is the matter of interest for the researchers. The anti-arthritis, antispasmodic, antibacterial, anti-inflammatory, immunostimulant, antidiabetic, hepatoprotective, antioxidant, antimicrobial, anthelmintic, antileishmanial, anti-pyretic, anti-allergic, antiviral and CNS depressant activities of the plant show its great value in the field of medicine. Considering NAT for treatment of various ailments can provide effective and efficient alternative against chemical drugs, which have no side effects and are cost-effective. Further attention and research are required for identification and characterisation of bioactive compound(s) responsible for the biological activity of plant and the elucidation of the mechanism of action in many cases. The toxicity of the various extracts should be considered, as human studies for safety and efficacy of the extracts for long term administration are needed to be proved. Use of various parts of NAT plants in Ayurveda, Siddha and Unani system of medicine should be considered, explored and effective treatment should be elucidated. Tissue culture, recombinant DNA technology and molecular marker-based approaches can be undertaken for increasing production, selection of desired traits of the plant and for studying diversity in germplasm of the species. Other chemical compounds like putrescine should be identified

that would help in increasing productivity. Combinational effect of various herbal plants along with NAT can be examined which could provide best alternatives for various ailments as was examined in case of wound healing activity of ethanolic extract of *Nyctanthes arbor-tristis* and *Murraya koenigii* by Jadhav *et al*, 2017. High potential of plant in management of various ailments, easy availability and requirement of no special condition for its collection and cultivation make it a plant of clinical interest which requires more attention and clinical trials for manufacturing therapeutic preparations that can treat human ailments.

### References

1. Smita Parekh, Anjali Soni, *Nyctanthes arbor-tristis*: Comprehensive review on its pharmacological, antioxidant and anticancer activities, *Journal of Applied Biology and Biotechnology*, 2020;8(01):95-104.
2. Anowar Hussain, Anand Ramteke, Flower extract of *Nyctanthes arbor-tristis* modulates glutathione level in hydrogen peroxide treated lymphocytes, *Pharmacognosy Res.*, 2012;4(4):230-233. DOI-10.4103/0974-8490.102272
3. Milind M Meshram, Swatee B Rangari, Shashank B Kshirsagar, Suraj Gajbhiye, Madhurim R Trivedi *et al*. *Nyctanthes arbor-tristis* a herbal panacea, *International Journal of Pharmaceutical Science and Research*, 2012, Page No: 2432-2440, DOI-[http://dx.doi.org/10.13040/IJPSR.0975-8323.3\(8\).2432-40](http://dx.doi.org/10.13040/IJPSR.0975-8323.3(8).2432-40)
4. K. Priya, Deepak Ganjewala, Antibacterial activities and Phytochemical analysis of different plant parts of

- Nyctanthes arbor-tristis (Linn.), Research Journal of Phytochemistry, 2007;1:61-67. DOI: 10.3923/rjphyto.2007.61.67
5. Laxmi Verma, Vaibhav Tamrakar, Narul Haque, Anil Kumar. Antifungal activity of different parts of Nyctanthes arbor-tristis Linn. (Parijat) against clinical pathogens, Shodh Darpan (Special Issue), 2016;1(4). ISSN No. 2454-1516
  6. Vidyavati Hiremath, Hiremath BS, Mohapatra S, Arun Kumar Das. Literary review of Parijata (Nyctanthus arbor-tristis Linn.) An herbal medicament with special reference to Ayurveda and Botanical literature, Biomed Pharmacol J, 2016, 9(3), DOI-<https://dx.doi.org/10.13005/bpj/1043>
  7. Reema Srivastava, Deepali Trivedi, Gauri Shukla, Pankaj Srivastava, Nyctanthus arbor-tristis: A wonder Indian herbal drug needs healthcare attention, Biomedical Journal of Scientific and technical research, published on June 11, 2018, ISSN: 2574-1241, DOI:10.26717/BJStr.2018.05.001199
  8. Champa Rani, Sunaina Chawla, Manisha Mangal, AK Mangal, Subhash Kajla, AK Dhawan *et al*, Nyctanthes arbor-tristis Linn. (Night Jasmine): A sacred ornamental plant with immense medicinal potential, Indian Journal of Traditional Knowledge, 2011;11(3):427-435.
  9. Jadhav VS, Ghawate VB. Evaluation of combined wound healing activity of ethanolic extracts of leaves of Murraya koenigii and Nyctanthes arbor-tristis on rats, Drug Invention Today, 2017;9:2. ISSN: 0975-7619
  10. Sasmal D, Sanjita Das, Basu SP. Review article on Phytoconstituents and therapeutic potential of Nyctanthes arbor-tristis Linn., Pharmacognosy reviews, July-Dec, 2007;1:2.
  11. Suresh V, Arunachalam G, Senthil Kumar N, In vitro anthelmintic activity of Nyctanthes arbor-tristis Linn. Bark, Journal of Pharmacy Research, 2011;4(1):283-284. ISSN: 0974-6943
  12. Bansal Gulshan, Suri KA, Grover Parul, A comprehensive review on Nyctanthes arbor-tristis, Int. J. Drug Dev. & Res., ISSN 0975-9344, January – March, 2015, 7.
  13. Ansari Intiyaz Ahmed, Patil Javesh K, Kamora Kunal Rajeshlal, Mandhan Priyanka Mahesh, Poptani Taruna Hareshlal, Girase Ananda Narayan. Anthelmintic activity of Nyctanthes arbor-tristis leaves on Indian earth worms, Indian Journal of Drugs, 2016;4(2):63-68. ISSN: 2348-1684
  14. Jadhav Santosh, Patil Manoj kumar. A Review on Nyctanthes arbor-tristis Linn. Rejuvenating herbs, International Journal of Research in Pharmacy and Pharmaceutical Science, 2016;1(1):54-62. March ISSN: 2455-698X
  15. Sonakshi Antal, Vrish Dhvaj Ashwlayan, Anuj Kumar. Protective effect of Nyctanthes arbor-tristis on scopolamine induced amnesia in rats “A Behavioural and Biochemical approach”, International Journal of Pharmaceutical Sciences and Research, Sr No. 48, Page No.: 1346-1355, Accepted on: 01 December’ 2016, published on: 01 March’ 2017, DOI: 10.12040/IJPSR.0975-8232.8(3).1346-55
  16. Mark S. Le Doux, Murine models of Caytaxin deficiency, in Movement Disorders (Second edition), pages, 2015, 439-452.
  17. Sunil A. Nirmal, Subodh C. Pal, Subhash C. Mandal, Antihistaminic activity of Nyctanthes arbor-tristis bark, article in Pharmacologyonline, 2009;3:924-928.
  18. Ramadan B. Sopi, Muhammad Feroz Hayat Khan, Bronchodilatory effect of ethanolic extract of the leaves of Nyctanthes arbor-tristis, Pharmacognosy Res, 2013;5(3):169-172. DOI: 10.4103/0974-8490.112422
  19. Matadeen Bharti RC Saxena, Omarao Singh Baghel, Rahul saxena, Apte KG. Wound healing activity of Nyctanthes arbor-tristis (Linn.), International Journal of Pharmaceutical sciences and research, Published on 01 October, 2011, 2694-2698. DOI: [https://dx.doi.org/10.13040/IJPSR.0975-8232.2\(10\).2694-98](https://dx.doi.org/10.13040/IJPSR.0975-8232.2(10).2694-98)
  20. Vaibhav Mishra, Astha Shukla, Sukanya Pandeti, Manoj Kumar Barthwal, Haushila Prasad Pandey, Gautam Palit *et al*. Arbotristoside-A and 7-O-trans-cinnamoyl-6 $\beta$ -hydroxyloganin isolated from Nyctanthes arbor-tristis possess anti-ulcerogenic and ulcer-healing properties, Phytomedicine: International Journal of Phytotherapy and Phytopharmacology, 2013. DOI: 10.1016/j.phymed.2013.04.010
  21. Bibhuti Bhusan Kakoti, Pares Pradhan, Sudarshan Borah, Kabita Mahato, Mritunjay Kumar. Analgesic and anti-inflammatory activities of the methanolic stem bark extract of Nyctanthes arbor-tristis Linn., Biomed Research International, vol.2013, 29 July 2013, Article ID: 826295, DOI: <https://doi.org/20.1155/2013/826295>
  22. Chaitali Pattanayak, Pratyay Pratim Datta, Analgesic activity of Nyctanthes arbor-tristis leaves in rodent, Journal of Interculture Ethnopharmacology, 2013;2(2):105-112. DOI: 10.5455/jice.20130312093742
  23. Gupta P, Bajaj SK, Chandra K, Singh KL, Tandon JS. Antiviral profile of Nyctanthes arbor-tristis L. against encephalitis causing viruses, Indian Journal of Experimental Biology, 2005;43:1156-1160.
  24. Bramanage Sachini Rangika, Pavithra Dilakshini Dayananda, Dinithi Champika Peiris, Hypoglycemic and hypolipidemic activities of aqueous extract of flowers from Nyctanthes arbor-tristis L. in male mice, BMC Complementary and Alternative medicine, 2015;15:289. 24 August 2015, DOI: 10.1186/s12906-015-0807-0
  25. Sneha J Anarthe, Swetha M, Spoorthisudha P, Ganga Raju M. Antioxidant and Antihyperlipidemic activity for methanolic extract of Nyctanthes arbor-tristis leaves, Indo American Journal of Pharmaceutical Sciences, 2016;3(12):1470-1481. DOI: <http://doi.org/10.5281/zenodo.221657>
  26. Maliha Uroos, Zaigham Abbas, Shumaila Sattar, Nigarish Umer, Arham Shabbir, Shafiq-ur-Rehman *et al*. Nyctanthes arbor-tristis ameliorated FCA-induced experimental arthritis: A comparative study among different extracts, Hindwani Evidence Based Complementary and Alternative Medicine, 2017. Article ID 4634853, 13 pages, DOI: <https://doi.org/10.1155/2017/4634853>
  27. Sanjita Das, Dinakar Sasmal, Saumya Priya basu. Antispasmodic and Anthelmintic activity of Nyctanthes arbor-tristis Linn, International Journal of pharmaceutical Sciences and Research, Page No.: 51-

- 55, Published on 01, 2010. DOI: [http://dx.doi.org/10.13040/IJPSR.0975-8232.1\(2\).51-55](http://dx.doi.org/10.13040/IJPSR.0975-8232.1(2).51-55)
28. Sanjita Das, D. Sasmal, S. P. Basu, Evaluation of CNS depressant activity of different plant parts of *Nyctanthes arbortristis* Linn., *Indian Journal of Pharmaceutical Sciences*,2008:70(6):803-806. DOI: 10.4103/0250-474X.49129
  29. Shyamali Singha L, Meenakshi Bawari, Manabendra Dutta Choudhary. Hepatoprotective and Antipyretic effect of bark of *Nyctanthes arbortristis* Linn., *International Journal of Pharmacy and Pharmaceutical Sciences*,2014:6. Suppl 2, ISSN: 0975-1491
  30. David M Arnoff, Eric G Neilson. Antipyretics: Mechanisms of action and clinical use in fever suppression, *The American Journal of Medicine*,2001:111(4):304-15. DOI: 10.1016/s0002-9343(01)00834-8
  31. Mrunal K Shirsat, Gupta SK, Bele D, Vaya R, Dwivedi, Goyal S *et al.* Antipyretic effect of whole plant extract of *Nyctanthes arbortristis* Linn., *Research and Review: A Journal of Pharmaceutical Science*,2011:2(1):7-10.
  32. Shalini Tripathi PK, Tripathi M, Vijay kumar, Ch V Rao, Singh PN. Anxiolytic activity of leaf extract of *Nyctanthes arbor-tristis* in experimental rats, *Pharmacologyonline*,2010:2:457-463.
  33. Awadhesh Kumar Mishra, Prabhat Upadhyay, Jyoti Dixit, Pradeep Kumar, Kavindra Nath Tiwari, Sunil Kumar Mishra *et al.* ameliorative activity of ethanolic flower extract of *Nyctanthes arbortristis* (L.) against scopolamine-induced amnesic effect and profiling of active compounds using gas chromatography-Mass spectrometry and ultra-performance liquid chromatography-quadrupole-time-of-flight mass spectrometry, *Pharmacognosy Magazine*,2018:14(59):596-604. DOI: 10.4130/pm.pm\_448\_18
  34. Sandhya Kumari TD, Sudha Madhuri TD, Singara Charya MA, Subba Rao K. Antioxidant and anticancer activities of *Nyctanthes arbor-tristis*, *Academic Sciences International Journal of Pharmacy and Pharmaceutical Sciences*,2012:4(4). Accepted on 11 August ISSN: 0975-1491
  35. Birendra Nath Karan, Tanushree Singha, Tanmoy Guria, Puspita Roy, TK Maity, Bikash Chandra Pal. Evaluation of anti-cancer activity of methanol extract of *Nyctanthes arbor-tristis* Linn., *Research Journal of Pharmaceutical, Biological and Chemical Sciences*,2017:8(5):412-420. ISSN: 0975-8585
  36. Mahipal S Shekhawat, Manokari M, Ravindran CP. An improved in vitro propagation protocol of *Nyctanthes arbor-tristis*, *Journal of herbs, shrubs and medicinal plants*, 2016. DOI: 10.1080/10496475.2016.1211211
  37. Bansal S, Bharti AJ, Bansal YK. Efficient in vitro regeneration of a medicinal plant Harsingar (*Nyctanthes arbor-tristis* L.), *Plant Tissue Culture and Biotech*,2012:22(2):137-142.
  38. H David Raja, DI Arockiasamy. In vitro propagation of *Nyctanthes arbor-tristis* L. *Indian Journal of Biotechnology*,2017:16:366-369.
  39. Ashok K Rohilla, Anil Sindhu, Anil K Poonia. Influence of putrescine on in vitro shoot morphogenesis from shoot-tip explants of *Nyctanthes arbor tristis* (L.): An important medicinal plant, *International Journal of*