



Effect of weed management strategies on the growth and yield performance of blackgram

Suseendran K, Nidheesh S, S Jawahar, C Kalaiyarasan, S Kalaisudarson, A P Srinivasaperumal

Department of Agronomy, Faculty of Agriculture, Annamalai University, Annamalai Nagar, Tamil Nadu, India

Abstract

Field experiment was conducted at farmer's field at Gudamalai village, Gangavallitaluk, Salem district during *Rabi* season 2019 to evaluate the performance of different pre emergence and early post emergence herbicides on weeds and productivity of blackgram under irrigated conditions. The experiment was laid out in randomized block design with three replications. The experiment consists of ten treatments T₁- Pendimethalin @ 1.0 kg ha⁻¹ on 3 DAS, T₂- PE Pendimethalin @ 1.0 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g/ha (Odyssey), T₃- PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS, T₄- PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + HW on 25 DAS, T₅- PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g/ha (Odyssey), T₆- PE Oxadiazon @ 250 g ha⁻¹ on 3 DAS + EPOE Imazethapyr @ 70 g ha⁻¹ (2-3 leaf stage of weeds), T₇- PE Oxadiazon @ 250 g ha⁻¹ on 3 DAS + EPOE Imazethapyr @ 70 g ha⁻¹ + Quizalofop-ethyl @ 50g/ha(2-3 leaf stage of weeds), T₈- EPOE Imazethapyr @ 70 g ha⁻¹ + Quizalofop-ethyl @ 50g/ha (2-3 leaf stage of weeds)- Tank mix, T₉- Hand Weeding twice @ 20 and 40 DAS, T₁₀- Unweeded check. Among the different weed control treatments imposed hand weeding twice at 20 and 40 DAS (T₉) registered the maximum growth and yield of irrigated blackgram due to reduced weed growth, weed dry matter production and reduction in nutrient depletion by weeds and increased nutrient uptake by the crop and the maximum growth (Plant height, leaf area index and Dry matter production) yield attributes (Number of pods per plant, Number of seeds per pod) and Yield (Grain yield kg ha⁻¹, Haulm yield kg ha⁻¹) of irrigated blackgram due to reduced weed growth, weed dry matter production and reduction in nutrient depletion by weeds and increased nutrient uptake by crop and this was followed by pre emergence application of Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g/ha (Odyssey) (T₅). The higher weed density, weed dry matter production, WCE, HET and minimum growth and yield and yield attributes was recorded under unweeded check (T₁₀).

Keywords: blackgram, herbicides, growth, yield and yield attributes

Introduction

Pulses are the major source of supplementary protein to daily diets and starchy food for a predominantly vegetarian population. Pulses are cheaper than meat and therefore often regarded as poor man's meat. Pulses contain on an average of 20-25 percent protein (Sooraj Chandra Pankaj *et al.*, 2017) [14]. Pulses are wonderful gift of nature with the unique ability of biological nitrogen fixation, deep root system, mobilization of insoluble soil nutrients and bringing qualitative changes in soil properties which make them known as soil fertility restorers (Deepak Kumar *et al.*, 2018) [5]. The total world acreage under pulses is about 85.40 million hectares with a production of 87.40 million tonnes at 1023 kg ha⁻¹ yield level. In India, pulses are being cultivated under an area of 29.36 million hectares with the production of 24.51 million tonnes and productivity of 835 kg ha⁻¹. The total area under pulses in Tamil Nadu is 8.16 lakh hectares with a production of 5.72 lakh tonnes. (DES, Ministry of Agriculture & FW (DAC & FW), Govt of India, (2018). Their productivity of pulses can be doubled by improved cultivars and by modern production technologies Sooraj Chandra Pankaj *et al.*, 2017) [14].

Blackgram (*Vignamungo* (L.)) is one of the most important pulse crops, which can be grown in tropical and subtropical regions. It is native of India and originated from *Phaseolus sublobatus* a wild plant. (Deepak Kumar *et al.* 2018) [5]. Blackgram is very nutritious as it contains a high level of carbohydrate (60g/100g), protein (20-25 g/100 g), phosphorus (385 mg/100 g), calcium (145 mg/100 g) and

iron (7.8 mg/100 g). It is useful in mitigating elevated cholesterol levels. The total world acreage under blackgram is about 23.48 million hectares with a production of 15.43 million tonnes at 653.07 kg ha⁻¹ yield level. The total area under blackgram in India is around 4.47 million hectares with a production of 2.83 million tonnes and productivity of 632 kg ha⁻¹. In Tamil Nadu, blackgram is cultivated in 4.30 lakh hectares with the production of 2.74 lakh tonnes and an average productivity of 637 kg ha⁻¹ (Indiastat, 2019) [8]. In blackgram there are various factors responsible for the reduction in growth and yield. The main constraint in blackgram production is weed infestation during its growth period which inflicts heavy losses on the crop yield by competing for essential growth factors. Weed management at the early stages of crop growth is essential, leading to severe competition between the crop and weeds. (Sukumar *et al.* 2018) [15]. Weeds compete for water, nutrient and space and cause up to 45% yield loss in Blackgram (Yadav *et al.*, 2015) [16]. An initial period of crop weed competition of 20-40 days is very critical and weed competition reduce blackgram yield to the extent of 87 percent. (Arvindverma *et al.* 2017). Weed management is an important key factor for enhancing the productivity of irrigated blackgram since the crop is not very good competitor against weeds (Choudhary *et al.* 2012) [4]. Hand weeding during critical crop growth stages is not possible due to increased cost and scarcity of human labour. Application of pre emergence herbicide lasts long only a very short period of the crop growth period. After the loss of

herbicide concentration weed seed bank start to emerge and compete with crops for natural resources. Delayed removal of weeds is not as effective in controlling weeds. Application of early post emergence herbicide controls late emerging weeds and obtains higher yields against the timely removal of weeds (Pratap Singh *et al.* 2016)^[12]. Under such circumstances, pre and early post emergence herbicides applied in sequence or combination will controls the weeds very effectively. Keeping these points in view, the present study was undertaken to find out the effect of pre and early post emergence herbicides and hand weeding practices on growth and yield of irrigated blackgram.

Materials and methods

The experiment was conducted at Gudamalai village, Gangavalli Taluk, Salem District, Tamil Nadu. The experimental farm is at 11°46' N and 78°59' E an altitude of 230 meters above MSL. The mean annual rainfall of the place is 952.2 mm. The mean maximum and minimum temperature prevailed during the cropping period was 34.9°C and 26.3°C respectively. The mean relative humidity ranges from 60.5 to 77.4 percent. The texture of the experimental field soil was sandy clay loam with neutral pH and low, medium and high in available nitrogen, phosphorus and potassium respectively. The popular variety Vamban (Bg) 5 was chosen for the study. The experiment was laid out in Randomized Block Design (RBD) with three replications and ten treatments. The treatment schedule were as follows : T₁- PE Pendimethalin @ 1.0 kg ha⁻¹ on 3 DAS, T₂ - PE Pendimethalin @ 1.0 kg ha⁻¹ on 3 DAS + EPOE

Imazethapyr + imazamox (RM) 70 g/ha (Odyssey), T₃- PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS, T₄ - PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + HW on 25 DAS, T₅ - PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g/ha (Odyssey), T₆ - PE Oxadiazon @ 250 g ha⁻¹ on 3 DAS + EPOE Imazethapyr @ 70 g ha⁻¹ (2-3 leaf stage of weeds), T₇ - PE Oxadiazon @ 250 g ha⁻¹ on 3 DAS + EPOE Imazethapyr @ 70 g ha⁻¹ + Quizalofop-ethyl @ 50g/ha (2-3 leaf stage of weeds), T₈ - EPOE Imazethapyr @ 70 g ha⁻¹ + Quizalofop-ethyl @ 50g/ha(2-3 leaf stage of weeds) – Tank mix, T₉ - Hand Weeding twice @ 20 and 40 DAS, T₁₀ - Unweeded check. The seeds were dibbled at the rate of 15 kg ha⁻¹ with the spacing of 30 × 10 cm. The pre emergence herbicides and early post emergence herbicides were applied on 3 DAS and 12 to 15 DAS as per the treatment schedule. Hand weeding was done on 20 and 40 DAS as per treatment schedule. Observations were recorded using quadrant (0.5 × 0.5 m). The biometric observations on Weed, growth and yield characters were recorded from five randomly selected plants.

Results and Discussion

Weed flora associated with irrigated blackgram

The major weed species of the experimental field comprised of grass, sedge and broad leaved weeds. The weed species are *Cynodon dactylon* and *Chloris barbata* among grass, *Cyperus rotundus* among sedge and *Parthenium hysterophorus*, *Phyllanthus niruri* and *Tridax procumbens* among broad leaved weeds.

Table 1: General weed flora of the experimental field

S.No	Botanical Name	Common Name	Habit	Family
A. Grass				
1	<i>Cynodon dactylon</i>	Bermuda grass	Perennial	Poaceae
2	<i>Chloris barbata</i>	Wind mill grass	Perennial	Poaceae
B. Sedge				
3	<i>Cyperus rotundus</i>	Nut sedge	Perennial	Cyperaceae
C. Broad leaved weeds				
4	<i>Parthenium hysterophorus</i>	Congress weed	Annual	Asteraceae
5	<i>Phyllanthus niruri</i>	Keezhanelli	Annual	Phyllanthaceae
6	<i>Tridax procumbens</i>	Tridax daisy	Annual	Asteraceae

Effect of weed management practices on weed Parameters

Weed density

The density of the total weed population in this experiment was significantly influenced by different weed management treatments in irrigated blackgram. The practice of hand weeding twice on 20 and 40 DAS (T₉) registered superior results over other treatments and recorded the least total weed population of 19.50 and 9.80 m⁻² on 30 and 45 DAS respectively. This was followed by pre emergence application of Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g ha⁻¹ (Odyssey) (T₅) and recorded the total weed population of 26.62 and 43.35 m⁻² on 30 and 45 DAS, respectively. These herbicides effectively controlled the broad spectrum of weeds like grass, sedges and broad leaved weeds. This might be due to the effective control of weeds throughout the season and persistent weed free condition for the rest of the crop growth period. This result corroborates with the findings of Balyan *et al.*, (2016)^[2]. Higher weed population was observed under unweeded check (T₁₀) with a total weed

population of 147.50 and 193.20 m⁻² on 30 and 45 DAS respectively.

Total weed dry matter production (kg ha⁻¹)

The total weed dry matter production was significantly reduced by weed control treatments. The least total weed dry matter production of 257.57 and 118.76 kg ha⁻¹ on 30 and 45 DAS was recorded in hand weeding twice on 20 and 40 DAS (T₉) which gave better results over other weed control treatments and this was followed by the herbicidal treatment PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g ha⁻¹ (Odyssey) (T₅) with weed dry matter production of 445.08 and 612.16 kg ha⁻¹ on 30 and 45 DAS respectively. The better performance of combination of herbicides might be due to synergistic effect between the two herbicides reducing the population as well as dry matter accumulation of different weed species. Similar results were indicated by Pratap Singh *et al.*, (2016)^[12]. Lesser weed DMP in this treatment was due to better control of weeds especially in the early stages of crop growth. Further, it was observed that

the lower dry matter production of weeds might be due to prolonged persistence of above herbicide contributing to weed free conditions during crop weed competition period (Patel *et al.*, 2015) [11]. Higher weed dry matter production was recorded under unweeded check (T₁₀) with total weed dry matter production of 1357.57 and 1607.12 kg ha⁻¹ on 30 and 45 DAS, respectively.

Weed control efficiency

The weed control efficiency in this experiment was significantly influenced by the different weed management practices in the irrigated blackgram. The highest weed control efficiency was observed on 30 and 45 DAS in hand weeding twice on 20 and 40 DAS (T₉) and was superior over others and recorded the highest weed control efficiency of 86.81 and 94.92 percent on 30 and 45 DAS and this was followed by the herbicidal treatment PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g ha⁻¹ (Odyssey) (T₅) with weed control efficiency of 81.95 and 77.56 percent on 30 and 45 DAS respectively, which reduced crop weed competition resulted in higher weed control efficiency, dry matter production and also removed grass, sedge and broad leaved weeds effectively in the critical stages of crop growth. Similar results were indicated by Sakthi, J. and A. Velayutham. 2018. Among the different weed control treatments, the lowest weed control efficiency of 61.15 and 60.80 percent was recorded in the treatment of PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS (T₃).

Herbicide Efficiency Index (HEI) (%)

The herbicide efficiency index was significantly influenced by the different weed management practices in the irrigated blackgram. The maximum highest herbicide efficiency index was observed in PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g ha⁻¹ (Odyssey) (T₅) and was superior over others and

recorded the highest herbicide efficiency index of 2.35 percent and this was followed by the herbicidal treatment PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + HW on 25 DAS (T₄) performed better in control of weeds throughout the cropping period among the herbicidal treatments. Maximum percentage of weeds eradicated in this treatment, which increases the higher Herbicide efficiency index (HEI). Similar findings were reported by Pratap Singh *et al.*, (2016) [12] with herbicide efficiency index of 1.57 percent. Among the different weed control treatments, the lowest herbicide efficiency index of 0.23 percent was recorded in the treatment of PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS (T₃).

Nutrient removal by weeds (kg ha⁻¹)

All the treatments exerted significant influence on nutrient removal by weeds in irrigated blackgram. The least N, P and K removal by weeds were recorded in hand weeding twice on 20 and 40 DAS (T₉) which gave better results over weed control treatments with the values of 24.86N, 10.24P and 18.72K kg ha⁻¹ respectively and this was followed by the herbicidal treatment PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g/ha (Odyssey) (T₅) with nutrient removal by weeds of 28.06N, 11.99P and 20.84K kg ha⁻¹ respectively. This might be due to efficient control of the emergence and growth of all weeds resulting in the least weed dry matter production and minimal nutrient removal by weeds. These findings correlate with the results of Khot *et al.*, (2016) [9]. The maximum nutrient removal by weeds was recorded under unweeded check (T₁₀) with the values of 42.83, 20.98, and 32.63 kg NPK ha⁻¹, respectively. The higher removal of nutrients in the weedy check might be attributed to higher weed count and greater dry matter accumulation by weeds. These findings were in accordance with the results of Hassan and Baswaid (2004) [7].

Table 2

Treatments	Dry Matter Production (Kgha ⁻¹)		Weed control efficiency (%)		Herbicide Efficiency Index	Nutrient Removal (Kgha ⁻¹)		
	30 DAS	45 DAS	30 DAS	45 DAS		N	P	K
T ₁ - PE Pendimethalin @ 1.0 kg ha ⁻¹ on 3 DAS	789.80	996.44	64.34	63.47	0.55	38.02	18.05	28.04
T ₂ - PE Pendimethalin @ 1.0 kg ha ⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g/ha (Odyssey)	696.89	872.73	71.74	68.01	1.18	33.58	15.52	24.88
T ₃ - PE Pendimethalin (38.7%) @ 0.75 kg ha ⁻¹ on 3 DAS	958.05	1067.47	61.15	60.80	0.23	40.38	19.47	30.29
T ₄ - PE Pendimethalin (38.7%) @ 0.75 kg ha ⁻¹ on 3 DAS + HW on 25 DAS	595.06	778.61	77.23	71.56	1.57	31.02	13.95	22.87
T ₅ - PE Pendimethalin (38.7%) @ 0.75 kg ha ⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g/ha (Odyssey)	445.08	612.16	81.95	77.56	2.35	28.06	11.99	20.84
T ₆ - PE Oxadiazon @ 250 g ha ⁻¹ on 3 DAS + EPOE Imazethapyr @ 70 g ha ⁻¹ (2-3 leaf stage of weeds)	740.36	902.55	69.98	66.91	0.90	34.93	16.21	25.65
T ₇ - PE Oxadiazon @ 250 g ha ⁻¹ on 3 DAS + EPOE Imazethapyr @ 70 g ha ⁻¹ + Quizalofop-ethyl @ 50g/ha(2-3 leaf stage of weeds)	719.92	888.14	70.81	67.44	1.06	34.22	15.48	25.29
T ₈ - EPOE Imazethapyr @ 70 g ha ⁻¹ + Quizalofop-ethyl @ 50g/ha(2-3 leaf stage of weeds) – Tank mix	840.18	979.02	65.93	64.13	0.65	37.22	17.74	27.68
T ₉ - Hand Weeding twice @ 20 and 40 DAS	275.57	118.76	86.81	94.92	-	24.86	10.24	18.72
T ₁₀ - Unweeded check	1357.57	1607.12	0	0	-	42.83	20.98	32.63
S.Ed	35.49	18.39	1.33	1.10	14.19	0.97	0.48	0.70
CD(P= 0.05)	71.69	37.15	2.69	2.23	28.67	1.95	0.96	1.42

Effect of weed control methods on growth and yield

Effect on growth attributes (Table: 1)

Among the weed control treatments, the plant height, leaf area index and DMP significantly influenced by various treatments. On 30 and 45 DAS higher plant height (20.42, 36.00 cm), leaf area index (4.95 on flowering), DMP (872, 2284 kg ha⁻¹) was registered in hand weeding twice on 20 and 40 DAS (T₉) and this was followed by the herbicidal treatment PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g ha⁻¹ (Odyssey) (T₅) with a plant height (18.89 and 33.48 cm) leaf area index (4.52 on flowering stage), DMP (840, 1928 kg ha⁻¹) on 30 and 45 DAS respectively. A weed free environment until the critical period of the crop growth by hand weeding facilitated good growth of the crop. Improved nutrient uptake and vigour due to elimination of weed competition right from the beginning of the crop might have contributed to favorable growth components, higher nutrient uptake and consequently higher plant height, LAI and DMP

in hand weeding twice treatment. The superiority of hand weeding practice at 20 and 40 DAS may be attributed to better weed control, least nutrient accumulation by weeds and better aeration of the crop. Similar findings were revealed by Manoj Kumar Sandil *et al.*, (2015)^[10]. Similarly, in herbicide application of Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g ha⁻¹ (Odyssey) (T₅) applied plots caused reduction in weed density, which might have improved the availability of resources viz., space, soil, moisture, light and nutrients to the irrigated blackgram, thus resulted in higher growth parameters. The increase in growth attributes under these treatments might be attributed due to the reduction in weed competitiveness with the crop, which ultimately favored better environment for growth and development of crop. Similar results are reported by Harithavarthini *et al.*, (2016). The least plant height, leaf area index, DMP was recorded under unweeded check (T₁₀).

Table 3: Influence of weed management practices on growth

Treatment	Plant Height (cm)		Leaf Area Index	Dry matter production at harvest (kg ha ⁻¹)	
	30 DAS	45 DAS		30 DAS	45 DAS
T1	12.68	24.28	2.92	664	1531
T2	15.82	28.59	3.69	773	1774
T3	11.20	21.79	2.51	627	1448
T4	17.38	31.01	4.1	809	1852
T5	18.89	33.48	4.52	840	1928
T6	14.98	27.38	3.43	726	1668
T7	15.41	27.95	3.57	749	1722
T8	13.20	24.87	3.03	687	1587
T9	20.42	36.00	4.95	872	2284
T10	9.62	19.27	2.10	585	1351
S.Ed	0.26	0.55	0.17	15.66	39.34
CD(P= 0.05)	0.52	1.12	0.34	31.64	79.47

Effect on yield attributes

All the treatments had a pronounced effect on the yield attributes of irrigated blackgram. Among the weed control treatments showed a noticeable influence on the number of pods plant⁻¹ on harvest (36.85) and pod length (5.5 cm) was recorded in hand weeding twice on 20 and 40 DAS (T₉) and this was followed by the herbicidal treatment PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g ha⁻¹ (Odyssey) (T₅) with the number of pods plant⁻¹ of 33.84 and pod length (5.1 cm) on harvest respectively. Among the treatments, hand weeding twice on 20 and 40 DAS (T₉) provided a perfect weed free environment throughout the critical period of crop growth and offered the highest value of yield components in the crop. This might be attributed to reduced crop weed competition in the critical stages which helped in synchronization of their production by increasing the number of pods plant⁻¹, number of seeds pod⁻¹ and pod length. These results were in agreement with the findings of Chhodavadia *et al.*, (2013)^[3], Manoj Kumar Sandil *et al.*, (2015)^[10] and Khot *et al.*, (2016)^[9]. Among the various chemical weed control methods, Pre emergence application of Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g ha⁻¹ (Odyssey) (T₅) recorded the highest values of yield components and the herbicide treatment pre emergence application of Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + HW on 25 DAS (T₄) was next in the order. It could be attributed to

significantly lower weed population, dry matter accumulation and higher weed control efficiency of weeds and also due to weed free environment provided by this treatment which might have increased the translocation of assimilates from source to sink and hence the yield attributes viz., number of pods plant⁻¹, number of seeds pod⁻¹ and pod length increase in this treatment. These results were in agreement with the findings of Patel *et al.*, (2015)^[11]. The lowest yield attributes were recorded under unweeded check (T₁₀). Severe weed competition exerted by weeds for the available resources throughout the crop growth might have reduced the yield components.

Effect on yield

Among the different treatments hand weeding twice @ 20 and 40 DAS (T₉) registered the maximum grain and haulm yield and it was 51.09 and 40.91 percent higher over unweeded check (T₁₀). The highest grain yield (912 kg ha⁻¹) and haulm yield (1459 Kgha⁻¹) was recorded in hand weeding twice on 20 and 40 DAS (T₉) and the application of herbicidal treatment PE Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g ha⁻¹ (Odyssey) (T₅) was next best and recorded 54.37 and 44.95 percent of grain yield (857 Kgha⁻¹) and haulm yield (1328 Kgha⁻¹) over unweeded check (T₁₀). This would be due to effective control of weeds which reduced the crop weed competition and increased yield of blackgram. These results were in agreement with the findings of Khot *et al.*, (2016)

^{19]}. This would be due to effective control of weeds which reduced the crop weed competition and increased yield of blackgram. The unweeded check showed the real depiction of the aggressive nature of weeds on the growth of irrigated

blackgram. The lowest grain (466 Kg ha^{-1}) and haulm yield (597 Kg ha^{-1}) were recorded in unweeded check (T₁₀). This might be due to the severe competition between crop and weed for different resources.

Table 4: Influence of weed management practices on yield parameters and yield

Treatment	No. of pods Plant ⁻¹	No ofseeds pod ⁻¹	Pod length (cm)	Test weight (g)	Grain yield (kg ha ⁻¹)	Haulm yield (kg ha ⁻¹)
T1	23.38	5.72	3.6	3.8	621	835
T2	29.11	6.08	4.3	4.3	758	1057
T3	20.91	5.64	3.2	3.6	540	693
T4	31.47	6.17	4.7	4.5	805	1182
T5	33.84	6.29	5.1	4.7	857	1328
T6	27.03	5.97	4.1	4.1	700	992
T7	28.01	6.03	4.2	4.2	733	1026
T8	24.58	5.88	3.7	3.9	652	880
T9	36.25	6.40	5.5	4.9	912	1459
T10	18.33	5.30	2.9	3.5	466	597
S.Ed	1.12	-	0.17	-	19	31
CD (P= 0.05)	2.27	NS	0.35	NS	38	62.43

Conclusion

Based on the above findings, hand weeding twice at 20 and 40 DAS (T₉) recorded the least weed population, weed dry matter production, nutrient removal by weeds, higher weed control efficiency and weed control index and favouring higher growth attributes, yield attributes, grain yield and haulm yield in irrigated blackgram and it was followed by pre emergence application of Pendimethalin (38.7%) @ 0.75 kg ha⁻¹ on 3 DAS + EPOE Imazethapyr + imazamox (RM) 70 g ha⁻¹ (Odyssey) (T₅). holds promise as an agronomically sound, ecologically safe and economically viable technology for enhancing the yield of irrigated blackgram.

References

- Arvind Verma, Roshan Choudhary, Choudary RS. Efficiency of imazethapyr and its ready mix on weed growth and yield of blackgram. Chem. Sci. Rev. Lett.,2017;6(24):2474-2477.
- Balyan JK, Choudhary RS, Kumpawat BS, Roshan Choudhary. Weed management in blackgram under rainfed conditions. Indian J. Weed Sci.,2016;48(2):173-177.
- Chhodavadia SK, Mathukiya RK, Dobariya VK. Pre and post-emergence herbicides for integrated weed management in summer greengram. Indian J. Weed Sci.,2013;45(2):137-139.
- Choudhary VK, Suresh KP, Bhagawati R. Integrated weed management in blackgram (*Vignamungo*) under mid hills of Arunachal Pradesh. Indian Journal of Agronomy,2012;57(4):382-385.
- Deepak Kumar, Singh RP, Somasundaram J, Vidya Simaiya, Shweta Jamra. Effect of foliar application of nutrients on growth and development of blackgram (*Vignamungo* (L.) Hepper) under rainfed Vertisols of Central India. IJCS,2018;6(1):609-613.
- Harithavardhini J, Jayalalitha K, Ashoka Raní Y, Krishnaveni B. Efficacy of post emergence herbicides on weed control efficiency, partitioning of drymatter and yield of blackgram (*Vignamungo* (L.) Hepper).Int. J. Food, Agric. Veterinary Sci.,2016;6(2):39-44.ISSN: 2277-2079.
- Hassan AA, Baswaid AS. The effect of seed rate application on the growth and yield of Blackgram. University of Aden, Yemen. J. Nat. Appli. Sciences,2004;(1):1-10.
- Indiastat. Online data base, 2019. In: <http://www.indiastat.com>
- Khot AB, Sagvekar VV, Muthal YC, Panchal VV, Dhonde MB. Effect on summer blackgram (*Phaseolusmungo* L.) to different sowing time and weed management practices with respect to yield, quality and nutrient uptake. Indian J. Weed Sci.,2016;38:57-62.
- Manoj kumar Sandil JK, Sharma Pratik Sanodiya. Effect of chemical weed management on productivity of soybean. Agric. Sust. Devp.,2015;2(1):1-5.
- Patel KR, Patel BD, Patel RB, Patel VJ, Darji VB. Bio-efficacy of herbicides against weeds in blackgram. Indian J. Weed Sci.,2015;47(1):78-81.
- Pratap Singh V, Tej Pratap Singh, Singh SP, Kumar A, Kavita Satyawali, Akshita Banga. Weed management in blackgram with pre-mix herbicides Indian J. Weed Sci.,2016;8(2):178-181.
- Sakthi J, Velayutham A. Effect of pre and post emergence herbicides on weeds and productivity of blackgram under irrigated conditions. Madras Agric, 2018;J.,105(1-3):16-19.
- Sooraj Chandra Pankaj, Pradeepkumar, Dewangan. Weed management in blackgram (*vignamungo* L.) and residual effect of herbicides on succeeding mustard (*brassica juncea* L.) crop. Int. J. Curr. Microbiol. App. Sci.,2017;6(11):865-881.
- Sukumar J, Pazhanivelan S, Kunjammal P. Effect of pre emergence and post emergence herbicides on weed control in irrigated blackgram. Journal of pharmacognosy and phytochemistry., SP,2018;1:3206-3209.
- Yadav KS, Dixit JP, Prajapati BL. Weed management effects on yield and economics of blackgram.Indian J. Weed Sci.,2015;47(T2):136-138.