



Determination of primary and secondary metabolites in ethnobotanical plants collected from Bharatpur district of Rajasthan

Renwati Dhidhariya, Rishikesh Meena*

Department of Botany, University of Rajasthan, Jaipur, Rajasthan, India

Abstract

Ethnobotany is the study of plants which are being used by tribal people to treat various diseases. In the present investigation, on the basis of tribal knowledge, we collected 10 ethnobotanical plants of various families (*Azadiracta indica* A. Juss, *Calotropis procera*, *Ricinus communis*, *Withania somnifera*, *Argemone maxicana*, *Oscimum sanctum*, *Hemidesmus indicus* L., *Abrus precatorius*, *Hygrophila auriculata*, and *Vitex negundo*). Plant parts used as traditional therapeutic purpose were selected for the study. Primary metabolites (TSS, proteins, free amino acids, and lipids) and secondary metabolites (total phenolic compounds, flavonoids, alkaloids, tannins, terpenoids, saponins) were determined quantitatively by using standard methods. Results revealed that these plant parts were rich in these metabolites. So, it can be concluded that the medicinal properties of these plant parts are due to presence of important phytochemicals.

Keywords: ethnobotany, tribal knowledge, therapeutic purpose, phytochemicals etc

Introduction

Ethnobiology is the study of biological features in relation to race, people, or cultural groups. The study of human populations' direct interactions with the plants and animals in their surroundings is known as ethnology.

J.W. Harshberger used the term "ethnobotany" in 1896 to describe the study of plants that were traditionally utilised by indigenous people. Stephen Power invented the phrase "aboriginal botany" in 1875 to characterise the study of all kinds of the vegetative world that the indigenous people of Australia employed for various purposes, including medicine, food, textiles, clothing, ornaments, and so on. Since Harshberger, the meaning of the term "ethnobotany" has undergone a process of change and development that is parallel to the formation and development of the science as a whole. For this reason, ethnobotany has been defined as the study of the relationship that exists between people from primitive societies and their plant environment (Schultes, 1962; Jain, 1976).

As a result of the presence of a large number of ethnic groups with a long history and a wide variety of plant life, India is one of the countries with the most extensive ethnobotanical knowledge. There are over 50 million people who belong to over 550 different tribal communities. The majority of these people live in the forest, hills, and other naturally remote areas. Ayurveda, often known as Ayurvedic medicine, is a form of complementary and alternative medicine that originated in India (Chopra and Ananda, 2003).

Plants that have been used medicinally for centuries produce a wide range of substances with established therapeutic uses (Ahmed and Beg, 2001). These substances are called as secondary plant metabolites. Primary metabolites serve as the building blocks for the biosynthesis of secondary metabolites, which encompass a vast variety of active chemicals. Their presence across the plant kingdom is far less widespread than that of other organisms. When grown in different environments, the same plant species can

produce varied amounts and qualities of these characteristics in varying degrees. They often accrue in lower quantities and have a tendency to be generated by particular cell types at various phases of development (Yeoman *et al.*, 1982; Endress, 1994; Edwards and Gatehouse, 1999). There are many plants secondary metabolites that are constitutive, meaning that they are present in healthy plants in their biologically active forms. Other plant secondary metabolites, on the other hand, occur as inactive precursors and are activated in response to tissue damage or the presence of pathogens (Osbourne, 1996).

In the present investigation, we investigated for determination of various primary and secondary metabolites in different ethnobotanical plants found in Bharatpur district of Rajasthan, India. In the district, about 12.44 percent of the population is comprised of people who belong to tribes such as the Bhil, Bhil-Meena, Damor, Dhanka, Garasia, Kathodi, Kokna, Kolidhor, Naikara, Patelia, Meena, and Seharla and who live in places that lack even the most fundamental infrastructure facilities. The cultural history of Rajasthan is further enriched by the presence of several nomadic tribes, including the Banjara, Gadolia-Lohar, Kalbelia, Sikligar, Kanjar, Sansi, and Bagri.

Materials and methods

Study area

The western Indian state of Rajasthan contains Bharatpur district as one of its many districts. The region known as Bharatpur District may be found in the eastern section of the state of Rajasthan. Its coordinates are as follows: 26°22' to 27°50' north latitude, 76.53' to 78.17' east longitude. The region covers an area of 5,066 km² in total. Which is equivalent to 1.48% of the district of Rajasthan's total land area.

Collection and processing of plants

Healthy parts of various plants (selected on the basis of traditional knowledge) were collected from various localities of the districts. Those were- leaves and stem bark

of *Azadiracta indica* A. Juss (family Maliaceae), leaves of *Calotropis procera* (family Apocynaceae), seeds of *Ricinus communis* (Family Euphorbeceae), roots of *Withania somnifera* (family Solanaceae), leaves and flowers of *Argemone maxicana* (family Papaveraceae), leaves of *Oscimum sanctum* (family Lamiaceae), roots of *Hemidesmus indicus* L. (Family Asclepiadaceae), leaves of *Abrus precatorius* (Fabaceae), seeds and roots of *Hygrophila auriculata* (family Acanthaceae), and leaves and roots of *Vitex negundo* (family Verbenaceae). The selected plant parts were cleaned, air-dried, grinded and stored in airtight boxes.

Determination of primary metabolites

Total soluble sugars

The total soluble carbohydrate content was determined according to the method described by Hedge and Hofreiter (1962). 1 ml of sample was mixed with 4 ml of anthrone reagent. Incubated in boiling water bath for 8 minutes after which the absorbance was read at 630 nm against a reagent blank. The analysis was performed in triplicates and the results were expressed as mg/g sample.

Proteins

Protein content was determined according to the method of Lowry *et al.* (1957). 1 ml of sample was mixed with 0.5 ml of 0.1 N NaOH and 5 ml of alkaline copper reagent, incubated the mixture in room temperature for 30 minutes. Added 0.5 ml of Folin-Ciocalteu reagent and incubated again for 10 minutes at room temperature. Absorbance was read at 660 nm against a reagent blank. The analysis was performed in triplicates and the results were expressed mg/g sample.

Determination of total free amino acids

Total free amino acids (ninhydrin method) was determined according to the procedure given by Moore and Stein (1948)^[13]. 1 ml of the sample was mixed with 1 ml of Ninhydrin in a test tube. Tubes were kept in boiling water bath for 20 minutes and then added 5 ml of diluent (equal volume of water and n-propanol) incubated at room temperature for 15 minutes and absorbance were read at 570 nm against a reagent blank. The analysis was performed in triplicates, and the results were expressed as mg/g sample.

Lipids

1gm of each dried peel sample was homogenised with 10 ml distilled water by using a mortar and pestle (Jayaraman, 1981). The paste obtained after homogenisation was mixed thoroughly with 30 ml of chloroform and methanol made in ratio of 2: 1 by v/v and transferred in a conical flask. Each mixture was left overnight at room temperature and then 20 ml of chloroform with equal volume of d.H₂O was added and centrifuged. After centrifugation three layers were obtained. Of these 3 layers, a transparent lower layer of chloroform containing all the lipids was poured out in pre-weighted beakers. Rest of the two layers were discarded. After total evaporation, the weight of beakers was measured again. The difference weight was considered as total lipids/g of the dried plant sample. The procedure was carried out in a set of three replicas and the mean value was noted for each sample.

Determination of secondary metabolites

Total phenolic compounds

The concentration of phenolics in plant extracts was determined using spectrophotometric method. Folin-Ciocalteu assay method was used for the determination of the total phenol content. The reaction mixture consists of 1 ml of extract and 9 ml of distilled water was taken in a volumetric flask (25 ml). One millilitre of Folin-Ciocalteu phenol reagent was treated to the mixture and shaken well. After 5 minutes, 10 ml of 7 % Sodium carbonate (Na₂CO₃) solution was treated to the mixture. The volume was made up to 25 ml. A set of standard solutions of gallic acid (20, 40, 40, 60, 80 and 100 µg/ml) were prepared in the same manner as described earlier. Incubated for 90 min at room temperature and the absorbance for test and standard solutions were determined against the reagent blank at 550 nm with an Ultraviolet (UV) /Visible spectrophotometer. Total phenol content was expressed as mg of GAE/gm of extract (Ghasemzadeh *et al.*, 2010; Rasool *et al.*, 2011)^[4].

Flavonoids

Total flavonoid content was measured by the aluminium chloride colorimetric assay. The reaction mixture consists of 1 ml of extract and 4 ml of distilled water was taken in a 10 ml volumetric flask. To the flask, 0.30 ml of 5 % sodium nitrite was treated and after 5 minutes, 0.3 ml of 10 % aluminium chloride was mixed. After 5 minutes, 2 ml of 1M Sodium hydroxide was treated and diluted to 10 ml with distilled water. A set of reference standard solutions of quercetin (20, 40, 60, 80 and 100 µg/ml) were prepared in the same manner as described earlier. The absorbance for test and standard solutions were determined against the reagent blank at 510 nm with an UV/Visible spectrophotometer. The total flavonoid content was expressed as mg of QE/g of extract (Kaviarasan *et al.*, 2007; Hanane *et al.*, 2010)^[9, 5].

Alkaloids

The plant extract (1mg) was dissolved in dimethyl sulphoxide (DMSO), added 1ml of 2 N HCl and filtered. This solution was transferred to a separating funnel, 5 ml of bromocresol green solution and 5 ml of phosphate buffer were added. The mixture was shaken with 1-4 ml chloroform by vigorous shaking and collected in a 10-ml volumetric flask and diluted to the volume with chloroform. A set of reference standard solutions of atropine (20, 40, 60, 80 and 100 µg/ml) were prepared in the same manner as described earlier. The absorbance for test and standard solutions were determined against the reagent blank at 470 nm with an UV/Visible spectrophotometer. The total alkaloid content was expressed as mg of AE/g of extract (Fazel *et al.*, 2008; Rao *et al.*, 2012)^[2, 19].

Tannins

The tannins were determined by Folin - Ciocalteu method. About 0.1 ml of the sample extract was added to a volumetric flask (10 ml) containing 7.5 ml of distilled water and 0.5 ml of Folin-Ciocalteu phenol reagent, 1 ml of 35 % Na₂CO₃ solution and dilute to 10 ml with distilled water. The mixture was shaken well and kept at room temperature for 30 min. A set of reference standard solutions of gallic acid (20, 40, 60, 80 and 100 µg/ml) were prepared in the same manner as described earlier. Absorbance for test and standard solutions were measured against the blank at 725 nm with an UV/Visible spectrophotometer. The tannin

content was expressed in terms of mg of GAE /g of extract . (Marinova *et al.*, 2005; Singh *et al.*, 2012; AfifyAel-M *et al.*, 2012) [12, 1].

Saponins

Saponin content was determined by the procedure of Obadoni and Ochuko (2001) [16]. The samples were ground and 20 g of each were put into a conical flask and 100 Cm³ of 20% aqueous ethanol were added. The samples were heated over a hot water bath for 4 h with continuous stirring at about 55°C. The mixture was filtered and the residue re-extracted with another 200 ml 20% ethanol. The combined extracts were reduced to 40 ml over water bath at about 90°C. The concentrate was transferred into a 250 ml separatory funnel and 20 ml of diethyl ether was added and shaken vigorously. The aqueous layer was recovered while the ether layer was discarded. The purification process was repeated and 60 ml of n butanol was added. The combined n-butanol extracts were washed twice with 10 ml of 5% aqueous sodium chloride. The remaining solution was heated in a water bath. After evaporation the samples were dried in the oven to a constant weight. The saponin content was calculated as percentage.

Percentage of Saponin = (Weight of residue /Weight of sample taken) × 100

Terpenoids

Terpenoids content was determined by the procedure of Ferguson, 1956. About 10 gm of leaves and bark powdered was taken and soaked in alcohol for 24 hours. It was filtered and filtrate extracted with petroleum ether; this ether extract was treated as total terpenoids (Kim *et al.*, 2003) [10].

Results and Discussion

Secondary metabolites are produced from primary metabolites in various metabolic pathways. These are stored in different plant parts as waste products. These secondary metabolites have various therapeutic uses for human beings. Traditional knowledge of using plants as therapeutic purpose is the basis of research on medicinal plants. On the basis on that traditional knowledge used by tribal people of Bharatpur district in Rajasthan, we selected some plant parts which are commonly used to treat various illnesses for determination of primary and secondary metabolites.

Results of quantitative analysis of various primary and secondary metabolites in the selected plants are shown in table 1 and table 2 respectively and these results are graphically represented in figure 1 and figure 2 respectively. Total soluble sugars (TSS), proteins, free amino acids, and lipids are primary metabolites which were determined in the present study. Results revealed that the used plant parts are rich in these primary metabolites which are precursors for various secondary metabolites. So, these plant parts were also found to be rich in secondary metabolites (total phenols, flavonoids, alkaloids, saponins, terpenoids, and tannins).

The selected plants are used being used to treat inflammation, hyperglycaemia, ulcers, malaria, micribial infections, infertility, diabetes, hypertension, cancer etc. (Subapriya and Nagini, 2005; Murti *et al.*, 2010; Prakash and Gupta, 2005) [20, 14, 17].

Results of the current study revealed the presence of important phytochemicals which are well known for showing medicinal properties (Jain *et al.*, 2019; Hussein and El-Anssary, 2018) [7]. Flavonoids are known to be natural antioxidants.

Table 1: Quantitative determination of primary metabolites in different plants collected from Bharatpur district of Rajasthan.

Name of plant	Part	Name of primary metabolites			
		TSS (mg/g.dw)	Proteins (mg/g.dw)	Free amino acids (mg/g.dw)	Lipids (mg/g.dw)
<i>Azadiracta indica</i> A. juss	Leaves	2.14	1.87	1.62	1.55
	Stem bark	1.03	0.64	1.38	1.78
<i>Calotropis procera</i>	Leaves	3.64	1.15	0.85	2.31
<i>Ricinus communis</i>	Seeds	1.17	1.32	1.25	2.86
<i>Withania somnifera</i>	Root	1.67	1.73	0.86	1.53
<i>Argemone maxicana</i>	Leaves	2.75	1.84	1.01	1.84
	Flowers	1.46	1.90	1.44	1.80
<i>Oscimum sanctum</i>	Leaves	2.93	1.35	0.78	1.54
<i>Hemidesmus indicus</i> L.	Roots	2.08	1.79	1.14	2.06
<i>Abrus precatorius</i>	Leaves	3.15	1.22	0.84	2.16
<i>Hygrophila auriculata</i>	Seed	1.73	1.08	1.36	3.72
	Root	2.15	1.57	0.72	2.26
<i>Vitex negundo</i>	Leaves	2.77	1.88	1.13	1.94
	Root	1.56	1.24	0.44	1.43

Table 2: Quantitative determination of secondary metabolites in different plants collected from Bharatpur district of Rajasthan.

Name of plant	Part	Name of Secondary metabolites					
		Total phenolic content (mg/g.dw)	Flavonoids (mg/g.dw)	Alkaloids (mg/g.dw)	Terpenoids (mg/g.dw)	Tannins (mg/g.dw)	Saponins (mg/g.dw)
<i>Azadiracta indica</i> A. juss	Leaves	1.79	1.06	0.68	0.88	1.15	0.73
	Stem bark	1.65	1.16	0.84	1.02	1.13	0.66
<i>Calotropis procera</i>	Leaves	1.26	0.96	0.68	0.34	0.17	0.57
<i>Ricinus communis</i>	Seeds	0.97	0.78	0.55	1.16	0.37	0.74
<i>Withania somnifera</i>	Root	1.66	1.28	0.87	0.65	0.53	0.08
<i>Argemone maxicana</i>	Leaves	1.09	0.76	0.54	0.47	0.83	0.26
	Flowers	1.23	0.88	0.23	0.68	0.57	0.14
<i>Oscimum sanctum</i>	Leaves	2.11	1.67	0.76	0.59	0.42	0.28

<i>Hemidesmus indicus</i> L.	Roots	0.87	0.68	0.52	0.16	0.23	0.73
<i>Abrus precatorius</i>	Leaves	1.35	0.85	0.44	0.18	0.72	0.38
<i>Hygrophila auriculata</i>	Seed	1.56	1.13	0.84	1.15	0.84	0.57
	Root	1.32	0.78	0.77	0.83	0.52	0.66
<i>Vitex negundo</i>	Leaves	1.40	1.21	1.17	0.75	0.26	0.88
	Root	1.28	1.06	1.25	0.68	0.47	0.61




Fig 1: Quantitative determination of primary metabolites in different plants collected from Bharapur district of Rajasthan.




Fig 2: Quantitative determination of secondary metabolites in different plants collected from Bharatpur district of Rajasthan.

Conclusion

From the results of the present investigation, it can be concluded that the selected plants are rich sources of pharmaceutically important phytochemicals. So, these can be used to identify and isolate active compounds for formulating medicines. Such natural medicines will be affordable with no/lesser side effects.

Acknowledgement

The authors are grateful to head, department of Botany, University of Rajasthan, Jaipur for providing facilities to conduct research work.

References

1. AfifyAel-M, El-Beltagi HS, El-Salam SM, Omran AA. Biochemical changes in phenols, flavonoids, tannins, vitamin E, β -carotene and antioxidant activity during soaking of three white sorghum varieties. Asian Pac J Trop Biomed.,2012;2(3):203-9.
2. Fazel S, Hamidreza M, Rouhollah G, Mohammadreza V. Spectrophotometric determination of total alkaloids in some Iranian medicinal plants. Thai J Pharm Sci.,2008;32:17-20.
3. Ferguson NM. A Text book of Pharmacognosy. Mac Milan Company, New Delhi, 1956, 191.

4. Ghasemzadeh AH, Hawa ZE, Asmah R. Antioxidant activities, total phenolics and flavonoids content in two varieties of Malaysia Young Ginger (*Zingiber officinale* Roscoe). Mol.,2010;15:4324-4333.
5. Hanane El, Hajaji, Nadya L, Katim A, Yahya C, Abdellah F *et al.* Antioxidant Properties and Total Phenolic Content of Three Varieties of Carob Tree Leaves from Morocco. Rec Nat Prod.,2010;4(4):193-204.
6. Hedge JE, Hofreiter BT, In: Whistler RL, Be Miller JN, editors. Carbohydrate Chemistry. New York: Academic Press,1962:17:1-19.
7. Hussein RA, El-Anssary AA. Plant secondary metabolites: the key drivers of the pharmacological actions of medicinal plants. Open access peer reviewed chapter, 2018.
8. Jain C, Khatana S, Vijayvergia R. Bioactivity of secondary metabolites of various plants: a review. International J of pharmaceutical sciences and research,2019;10(2):494.
9. Kaviarasan S, Naik GH, Gangabhagirathi R, Anuradha CV, Priyadarsini KI. *In vitro* studies on antiradical and antioxidant activities of fenugreek (*Trigonella foenumgraecum*) seeds. Food Chem.,2007;103:31-37.
10. Kim D, Jeong SW, Lee CY. Antioxidant capacity of phenolic phytochemicals from various cultivars of pulms. Food Chemistry,2003;81(3):321-326.
11. Lowry OH, Roseobrough NJ, Farr AL, Randall RJ. Protein measurements with the Folin's phenol reagent. J Biol Chem.,1957;193:265-75.
12. Marinova D, Ribarova F, Atanassova M. Total Phenolics and Total Flavonoids in Bulgarian Fruits and Vegetables. J University ChemTechnol Metallurgy,2005;40(3):255-260.
13. Moore S, Stein WH. Photometric methods for use in the chromatography of amino acids. J Bio Chem.,1948;176:367-88.
14. Murti Y, Yogi B, Pathak D. Pharmacognostic standardization of leaves of *Calotropis procera* (Ait.) R. Br. (Asclepiadaceae). Int J Ayurveda Res.,2010;1(1):14-17.
15. Nasir R, Komal R, Muhammad Z, Kaleem URN, Imran I, Viqar UA. Antioxidant potential of different extracts and fractions of *Catharanthusroseus* shoots. Int J Phytomed.,2011;1(3):108-114.
16. Obdoni BO, Ochuko PO. Phytochemical studies and comparative efficacy of the crude extracts of some Homostatic plants in Edo and Delta States of Nigeria. Global Journal Pure Appl. Sci.,2001;8b:203-208.
17. Prakash P, Gupta N. Therapeutic uses of *Ocimum sanctum* Linn (Tulsi) with a note on eugenol and its pharmacological actions: a short review. Indian J Physiol Pharmacol.,2005;49(2):125-31.
18. Rajeev S, Pawan KV, Gagandeep S. Total phenolic, flavonoids and tannin contents in different extracts of *Artemisia absinthium*. J Intercult Ethnopharmacol.,2012;1(2):101-104.
19. Rao MT, Rao BG, Rao YB. Antioxidant activity of *Spilanthes acmella* extracts. Int J Phytopharmacol.,2012;3(2): 216-220.
20. Subapriya R, Nagini S. Medicinal properties of neem leaves: a review. Curr Med Chem anticancer agents,2005;5(2):149-156.