



## A recent review on phytochemicals commonly found in herbal plants

T Sivakumar\*

Department of Botany, Annamalai University, Annamalai Nagar, Tamil Nadu, India

### Abstract

In the current review article novel discovery of traditional plants and herbal products are gaining famous in the scientific community present days owing to their reliable pharmacological properties and affordability to prices for the common people which develops them efficacy in control of various diseases. They have made significant contributions to rural livelihoods, and many, in addition to traditional healers who practice herbal medicine People are involved in collecting and trading medicinal plants. Thus, the global demand for herbal medicines has increased, which has led to the realization of advanced new medicines. Most communities in developing countries are deeply involved in the use of medicinal plants due to the alarming poverty rates, so this requires scientific support, especially in reducing their effectiveness in treating such diseases. In their pure compounds or standardized plant extracts, they offer unlimited opportunities for new drugs because they cannot compensate for their chemical diversity. The key of natural products is associated to the efficacy of different phytochemicals such as tannins, terpenoids, cardiac glycosides, flavonoids, saponins and alkaloids are treat as cancer activity. The available reports on the phytochemicals within the general herbal plants are discussed in this review.

**Keywords:** alkaloids, flavonoids, saponins, tannins, terpenoids

### Introduction

Plants are naturally gifted with a set of medicinal compounds whose characteristic has led to the discovery of new, cheaper drugs with greater therapeutic potential<sup>[1, 2]</sup>. Their exploitation in medicine is very ancient, for example, in 28 AD the Greek physician Dioscorides wrote *de Materia medica* containing 600 medicinal plants and was the leading text on pharmacology until the Renaissance. Hippocrates may have suggested the leaves of the willow tree to reduce fever. Salicylic acid, which has anti-inflammatory and analgesic properties, was first extracted from white willow and then synthetically made into a staple<sup>[3, 1]</sup>. The pharmacological resources of the herbal utilized for food, medicine or spiritual purposes for centuries have been confirmed by new approaches to their analysis<sup>[4-10]</sup>. The folk use of few plants in traditional medicine has been consistently confirmed by scientific research based on their effects<sup>[11-15]</sup>. The traditional use of certain plants in traditional medicine has been consistently confirmed by scientific research based on their effects<sup>[11]</sup>. This shift to herbal drugs has been facilitated by factors like the low cost of herbal drugs, endearing them with the poor mass of developing world; the 'green' movement in the developed world that campaigns on the inherent safety and desirability of natural products and the individualistic philosophy of western society that encourages self-medication, with many people preferring to treat themselves with phytomedicines<sup>[3]</sup>.

### Phytochemicals

Phytochemicals contains antioxidant effects, modulation of enzyme activity, induced of the immune system, antibacterial, hormonal metabolism and antiviral effects, DNA replication and / or body interaction. It can physically bind to certain cell walls, thereby preventing disease from attaching to human cell walls<sup>[16, 17]</sup>. The phytochemicals are explained in the following sections.

### Alkaloids

An alkaloid is a compound obtained from a plant Toxic or physiologically active. Some alkaloids contains isopteroprotein, terophobine, etc. The antimicrobial activity they perform Promotes the removal of harmful white blood cells Microbes and cell debris<sup>[18]</sup>. They act as narcotics and antimalaria. As a topical anesthetic for ophthalmology; In prolonging the life of hypertension, neurology, rheumatism, motor disease and hormones<sup>[17]</sup>. They have analgesic action, so it is used to reduce the pain of boils, septic injuries and complaints such as headaches, abdominal pain and eye conditions. They also have antinobiotic activity, for example indole alkaloids are used in leukemia and Hodgkin's disease chemotherapy. They act by stopping and depolymerizing the protein microtubules that make up the mitotic cycle in the cell division. This process helps to stop the tumor cells from dividing or dividing and as a result reduces the risk of cancer. However, Some types of alkaloids are illusory, Addictive, and poisonous and therefore arrow is used Poison for hunting wild game<sup>[18]</sup>.

### Saponin

Raising the respiratory in anticipations, and also it contains biological role, including anti-cough activity. They act as vaccine boosters by acting as adjuncts. They have anti-inflammatory, emetics, antiviral, antifungal, molluscicidal, piscidal and antibacterial activity<sup>[17]</sup>. The mechanism of action for antibacterial effects includes the membrane properties of saponin and the reduction of the surface tension of the extracellular medium<sup>[19]</sup>.

### Flavonoids

They are fundamental derivatives of flavonoids, have integrated aromatic systems, are often bound to glycosides with sugar(s), and are naturally phenolic and water-soluble<sup>[20]</sup>. They play their roles as antioxidants, thus protecting against degenerative pathogens. Flavonoids such as

quercetin act as chain-breaking antioxidants and inhibit the oxidation of low-density lipoprotein by metal ions such as macrophages and copper. It reduces antioxidant stress [17]. They act as antibiotics by complicating extracellular and soluble proteins and complicating the bacterial cell wall. Lipophilic flavonoids can also disrupt microbial membranes [21-23]. Potential targets of the microbial cell are surface-exposed adhesins, cell wall polypeptides, and membrane-bound enzymes. The catechins found in oolong green tea inactivate bacterial toxins (e.g. cholera toxins) and inhibit bacterial glucosyl transferase. They function; 'Nature's biological modifiers' trigger two enzymes that are anti-allergic, anti-inflammatory, and eliminate mutations and cancers [24]. They are also known to simplify capillary weakness [20], antiallergic and antispasmodic and therefore used for relief asthma and nasal bleeding [17]. Flavonoids that do not have hydroxyl groups (-OH) in their composition are more active against microorganisms than -OH, and this supports the notion of their microbial target membrane [25, 22, 17].

### Tannins

Tannins are astringent, bitter plant polyphenols that bind, accelerate or contract proteins. They play a physiological role by acting as antioxidants through free radical scavenging activity, transformation of metals, and fraud by proximate oxidative enzymes and lipid peroxidation [26-32, 17]. Modifying antioxidant pressure and preventing degenerative diseases. They inhibit tumor growth by inducing apoptosis [33] and by preventing the mutagenicity of cancers [34, 17]. They indicate antimicrobial activity by complicating nucleophilic proteins by hydrogen binding, covalent binding, and unspecified interactions. The chief targets of the complex are cell wall and cell membrane adhesive proteins, thus inactivating microbial adhesions, which is the first step in establishing infection. They also cause cell wall/membrane disorders [25, 34, 35, 17]. It also inactivates microenzymes and cell envelope transport proteins by processes involving reactions with sulfide groups of proteins [22, 23, 17].

They accumulate/ complexes metal ions (e.g. manganese, cobalt, copper, iron, etc.) required for microbial growth as co-agents and enzyme activators. They also inhibit viral reverse transcriptase [34, 35, 24, 17]. Toxicity to microorganisms in phenolic compounds depends on the site and number of hydroxyl groups, with evidence that hydroxylation leads to increased toxicity [36, 29]. They play an endocrine role by interacting with estrogen receptors. They are important in the control of anti-inflammatory, molluscicidal and schistosomiasis.

### Phytosteroids

Phytosteroids are plant steroids that may or may not act as weak hormones in the body. They share a common basic ring structure with animal stimulants, but they are not identical due to the various chemical groups attached to the main ring at different stages [17]. They are mainly used to treat reproductive problems, which are used to ensure easy delivery during pregnancy, as well as improve fertility in women and libido in men. They also act as sex hormone derivatives (for example, they can be metabolized to substances such as androgen or estrogen) so they are potential sources of contraception [37, 17]. They are utilized in antimicrobial, analgesic, anti-inflammatory and therapeutic

purposes and in reducing serum cholesterol levels [17]. They have also been revealed to be potent inhibitors of macrophage activity, inhibiting the production of anti-inflammatory cytokines and LPS-induced death, so they are particularly effective as prophylaxis agents especially the physalins [38-41].

### Terpenoids

These are the derivatives of the isoprene molecule with a carbon skeleton constructed from one or more C15 units [20]. They play their roles as antiviral, antibacterial, antifungal, protozoan, immune boosters, anti-allergic and antinoplastic [42]. The mechanisms of action by which these lipophilic compounds cause membrane degeneration are speculated [25, 24, 22]. Petalostemum has exhibit activity against *S. aureus*, *B. subtilis*, and *C. albicans* and to a lesser extent to Gram-negative bacteria [25, 17]. This may be due to a disruption of the lipid fraction of the bacterial plasma membranes, resulting in changes in membrane permeability and leakage of intracellular substances. This is associated to the physicochemical properties of the dynamic principle (such as lipophilicity and water solubility), lipid composition, and net surface charge of bacterial membranes. These phytochemicals can penetrate cell membranes, penetrate the cell's interior, and interact with important inward targets for antibacterial activity [43-52]. They are used to relieve epilepsy and to treat colds, coughs and acute bronchitis [17]. From laboratory studies of terbenes from ginseng, it is suggested that the potential target of these compounds includes the hypothalamus-pituitary-adrenal axis owing to the observed effects on adrenocorticotrophic hormone and corticosterone [52, 17].

### Conclusion

Herbal plant products are still an important plant in supporting primary health care systems. Some of these phytochemicals accidentally protect humans against pathogens, which is why they are a key target for drug anticipation programs. Their biological activity is mostly related to secondary metabolism, which is often described in detail for plant protection. The primary phytochemicals are known to have many properties that are important to cells, therapeutic, Prophylactic, nutritious and immune-modulating properties.

### Reference

1. Huie CW. Review: Modern sample preparation techniques for the extraction and analysis of medicinal plants. *Journal of Analytical and Bioanalytical Chemistry*, 2002:373:23-30.
2. Ukwuani AN, Abubakar MG, Hassan SW, Agaie BM. Antinociceptive Activity of Hydromethanolic Extract of Some Medicinal Plants in Mice. *International Journal of Pharmacy*. Photon, 2013:104:120-125.
3. Houghton PJ, Raman A. *Laboratory Hand book for the Fractionation of Natural Extracts*. London: Chapman and Hall, 1st edition, 1998, 1-153.
4. Mishra AP, Saklani S, Salehi B, Parcha V, Sharifi-Rad M, Milella L *et al.* a high altitude medicinal orchid of Indian Himalayan region: Chemical profile and biological activities of tuber extracts. *Cell and Molecular Biology*, 2018a:64:35-43.
5. Sharifi-Rad M, Ozcelik B, Altın G, Da,skaya-Dikmen C, Martorell M, Ramírez-Alarcón K *et al.* Alves Borges

- Leal AL. *Salvia* spp. plants-from farm to food applications and phytopharmacotherapy. *Trends Food Science and Technology*,2018;80:242-263.
6. Abdolshahi A, Naybandi-Atashi S, Heydari-Majd M, Salehi B, Kobarfard F, Ayatollahi SA *et al.* Antibacterial activity of some Lamiaceae species against *Staphylococcus aureus* in yoghurt-based drink (Doogh). *Cell and Molecular Biology*, (Noisy-le-grand),2018;64:71-77.
  7. Sivakumar T. Phytochemical screening and GC- MS analysis of bioactive compounds and biosynthesis of silver nanoparticles using sprout extracts of *vigna radiate* L. and their antioxidant and antibacterial activity. *Asian Journal of Pharmaceutical and Clinical Research*,2019;12(2):180-184.
  8. Deepa B. Sivakumar T. Phytochemical Analysis and Antibacterial Efficacy of Ethanolic Extract of *Musa paradisiaca*. *Research Biotica International Journal*,2020;2(3):126-130.
  9. Thiagarajan G, Sivakumar T. A Review on Pepper and Their Common Phytochemicals. *Research Biotica International Journal*,2020;2(4):149-153.
  10. Sivakumar T. *In vitro* antioxidant, total phenolic, total flavonoid content and biosynthesis of silver nanoparticles using *Cassia auriculata* leaves extracts. *International Journal of Botany Studies*,2021;6(3):476-480.
  11. Mishra AP, Sharifi-Rad M, Shariati MA, Mabkhot YN, Al-Showiman SS, Rauf A *et al.* Bioactive compounds and health benefits of edible *Rumex* species—A review. *Cell and Molecular Biology*,2018b;64:27-34.
  12. Angelin JJ, Jothi U, Thiagarajan G, Sivakumar T. Evaluation of antimicrobial activity and phytochemicals analysis of whole plant extract of *Vinca rosea*. *Asian Journal of Pharmaceutical and Clinical Research*,2019;12(8):132-136.
  13. Angelin JJ, Jothi U, Thiagarajan G, Subramanian VK, Sivakumar T. Estimation of qualitative and quantitative analysis of antioxidant activity of different parts of *Catharanthus roseus* (L). *Plant Archives*,2020;20(2):4187-4192.
  14. Sivakumar T, Panneerselvam R. Salinity induced changes in photosynthetic pigment and antioxidant responses in *Sesuvium portulacastrum*. *Pakistan Journal of Biological sciences*,2011;14(21):967-975.
  15. Jothi U, Angelin JJ, Sivakumar T. Study on Estimation and Antioxidant activity of *Gloriosasuperba* L. Whole Plant Extract. *International Journal Scientific Research and Biological Science*,2019;6(3):55-58.
  16. Abu-Darwish MS, Efferth T. Medicinal plants from near east for cancer therapy. *Frontiers in pharmacology*,2018;9:56.
  17. Ngoci SN, Mwendia CM, Mwaniki CG. Phytochemical and cytotoxicity testing of *Indigofera lupatana* Baker. *Frontiers Journal of Animal and Plant Sciences*,2011;11(1):1364-1373.
  18. Ogunwenmo KO, Idowu OA, Innocent C, Esan EB, Oyelana OA. Cultivars of *Codiaeum variegatum* (L.) Blume (Euphorbiaceae) show variability in phytochemical and cytological characteristics. *Journal of Biotechnology*,2007;20:2400-2405.
  19. Al-Bayati FA, Al-Mola HF. Antibacterial and antifungal activities of different parts of *Tribulus terrestris* L. growing in Iraq. *Journal Zhejiang University Science*,2008;9:154-159.
  20. Harborne JB. *Phytochemical methods*. London: Chapman and Hall, 1973, 52-114.
  21. Navarro MC, Montilla MP, Cabo MM, Galisteo M, Caceres A, Morales C, Berger I. Antibacterial, antiprotozoal and antioxidant activity of five plants used in izabal for infectious diseases. *Journal of Phytotherapy Research*,2003;17:325-329.
  22. Samy RP, Gopalakrishnakone P. Review: Therapeutic potential of plants as anti-microbials for drug discovery. *Evidence-Based Complementary and Alternative Medicine*, 2008. doi:10.1093/ecam/ nen036.
  23. Kaur GJ, Arora DS. Antibacterial and phytochemical screening of *Anethum graveolens*, *Foeniculum vulgare* and *Trachyspermum ammi*. *Journal of BioMedicine Central: Complementary and Alternative Medicine*,2009;9:30-4.
  24. Ogunwenmo KO, Idowu OA, Innocent C, Esan EB, Oyelana OA. Cultivars of *Codiaeum variegatum* (L.) Blume (Euphorbiaceae) show variability in phytochemical and cytological characteristics. *Journal of Biotechnology*,2007;20:2400-2405.
  25. Cowan MM. Plant products as anti-microbial agents. *Clinical Microbiology Reviews*,1999;12:564-582.
  26. Navarro MC, Montilla MP, Cabo MM, Galisteo M, Caceres A, Morales C *et al.* Antibacterial, antiprotozoal and antioxidant activity of five plants used in izabal for infectious diseases. *Journal of Phytotherapy Research*,2003;17:325-329.
  27. Sivakumar T. Synthesis of silver nanoparticles using *Cassia auriculata* leaf extracts and their potential antidiabetic activity. *International Journal of Botany Studies*,2021;6(3):35-38.
  28. Sivakumar T. A modern review of silver nanoparticles mediated plant extracts and its potential bioapplications. *International Journal of Botany Studies*,2021;6(3):170-175.
  29. Sivakumar T, Panneerselvam R. Triadimefon Mediated Changes in Antioxidant and Indole Alkaloid Content in Two Species of *Datura*. *American Journal of Plant Physiology*,2011;6: 252-260.
  30. Sivakumar T. A review on biosynthesis of Zinc Oxide Nanoparticles using plant extracts and its potential Bio Applications. *International Journal of Botany Studies*,2021;6(3):644-648.
  31. Sivakumar T, Deepa B. A Review on Some Folk Medicinal Plants and Their Common Uses. *Research Biotica International Journal*,2020;2(3):126-130.
  32. Shanmugapriya AK, Sivakumar T, Panneerselvam R. Study of antioxidant potentiality of tomato under tricyclazole and difenoconazole treatments. *International journal of Research in plant science*,2013;3(3):47-53.
  33. Scalbert A, Johnson IT, Saltmarsh M. Polyphenols: Antioxidants and beyond. *American Journal of Clinical Nutrition*,2005;81:215-217.
  34. Okuda T. Systematics and health effects of chemically distinct tannins in medicinal plants. *Journal of Phytochemistry*,2005;66:2012-2031.
  35. Biradar YS, Jagatap S, Khandelwal KR, Singhanian SS. Exploring of Antimicrobial Activity of *Triphala Mashi* - an Ayurvedic Formulation. *Journal of Evidence based*

- Complementary and Alternative Medicine,2007:5:107-113.
36. Przybylski R, Lee RY, Eskin NAM. Antioxidant and radical-scavenging activities of buckwheat seed components. Journal of the American Oil Chemists' Society,1998:75:1595-1601.
  37. Edeoga HO, Okwu DE, Mbaebie BO. Phytochemical constituents of some Nigerian medicinal plants. African Journal of Biotechnology,2005:4:685-688.
  38. Soares MB, Brustolim D, Santos LA, Bellintani MC, Paiva FP, Ribeiro YM *et al.* Physalins B, F and G, Seco lymphocyte function and allogeneic transplant rejection. Journal of International Immuno pharmacology,2005:6:408-414.
  39. Sivakumar T. A systematic review on traditional medicinal plants used for traditional control of insects. International Journal of Botany Studies,2022:7(12): 32-36.
  40. SenthilKumar SR, Sivakumar T, Arulmozhi KT, Mythili N. Antimicrobial Activity of Indian Commercial Green Teas (*Camellia Sinensis*). International Journal Bioscience and Nanoscience,2016:3(7):108-112.
  41. Senthil Kumar SR, Sivakumar T, Arulmozhi KT, Mythili N. FT-IR analysis and correlation studies on the antioxidant activity, total phenolics and total flavonoids of Indian commercial teas (*Camellia sinensis* L.) - A novel approach. International Research Journal of Biological Science,2017:6(3):1-7.
  42. Roberts SC. Production and engineering of terpenoids in plant cell culture. Journal of Nature Chemical Biology,2007:3: 387-395.
  43. Trombetta D, Castelli F, Sarpietro M, Venuti V, Cristani M, Daniele C *et al.* Mechanisms of anti-bacterial action of three monoterpenes. Journal of Antimicrobial Agents and Chemotherapy,2005:49:2474-2478.
  44. Senthilkumar SR, Sivakumar T. Green tea (*Camellia sinensis*) mediated synthesis of zinc oxide (ZNO) nanoparticles and studies on their antimicrobial activities. International Journal of Pharmacy and Pharmaceutical Sciences,2014:6(6):461-465.
  45. Senthilkumar SR, Sivakumar T. Studies on the Greengram (*Vigna radiata* L.) Sprout Assisted Synthesis of Silver Nanoparticles and their Antimicrobial Activities International Journal of Nanomaterial and Biostructure,2014:4(3):52-57.
  46. Senthilkumar SR, Sivakumar T, Arulmozhi KT, Mythili N. Gas chromatography Mass spectroscopy evaluation of bioactive phytochemicals of commercial green teas (*Camellia sinensis*) of India. Asian Journal of Pharmaceutical and Clinical Research,2015:8(3):278-282.
  47. Anburaj R, Nabeel M. Alikunhi Sivakumar T, Kathiresan K. Role of rhizobacteria on biochemical constituents of the mangrove associate halophyte *sesuvium portulacastrum* L., in response to salinity. Russian Journal of Plant Physiology,2012:59(1):115-119.
  48. Sivakumar T, Lakshmanan GMA, Murali PV, Panneerselvam R. Alteration of Antioxidative Metabolism Induced by Triazoles in Sweet Potato. Journal of Experimental Sciences,2010:(1)3:10-13.
  49. Sivakumar T, Sundaramanickam A, Panneerselvam R. Changes in growth and pigment content in sweet potato by Triadimefon and Hexaconazole. Journal of Phytology,2009:1: 333-341.
  50. Shanmugapriya AK, Sivakumar R, Panneerselvam R. Difenconazole and tricyclazole induced changes in photosynthetic pigments of *Lycopersicon esculentum*. L. International journal of Agriculture and food science,2013:3(2):72-75.
  51. Shanmugapriya AK, Sivakumar T. Regeneration of *in vitro* plantlets in *Hemidesmus indicus* (L.) R. Br. Through nodal and leaf explants. International multidisciplinary Research Journal,2011:1(10):41-45.
  52. Briskin DP. Medicinal plants and phytomedicines. Linking plant Biochemistry and Physiology to human health. Journal of Plant Physiology,2000:124:507-514.