

Summer pollen sources to *Apis dorsata* honeybees collected from Chandrapur tahsil forest area of Chandrapur district of Maharashtra state (India)

Borkar Laxmikant¹, Nakade Ankit², Mate Devendra²

¹ Department of Botany, S. S. Jaiswal Arts, Comm. and Science College, Arjuni (Mor), Gondia, Maharashtra, India

² Department of Botany, Nutan Adrash Arts, Commerce and M. H. Wegad Science College, Umrer, Nagpur, Maharashtra India

Abstract

76 pollen loads recovered directly from the honeycombs of *Apis dorsata* (Rock Bee) collected in 16 March 2012 to 3 May 2013 from Lohara and Chichpali forest area of Chandrapur Tahsil of Chandrapur District of Maharashtra State, were analysed. 8 (10.52%) pollen loads were found to be unifloral, 23 (30.26%) bifloral and 45 (59.21%) multifloral. The unifloral pollen loads were contained *Terminalia* sp. and *Mangifera indica*. The pollen of *Terminalia* sp. was recovered from 66 (86.84%) of the total pollen loads studied. The study highlights *Terminalia* sp. (Combretaceae), *Blumea* sp. (Asteraceae), *Mangifera indica* (Anacardiaceae), *Delonix regia* (Caesalpiniaceae), *Bombax ceiba* (Bombiaceae) as fairly important sources of pollen of the honeybees during the summer period.

Keywords: Pollen sources, honeybee, Chandrapur tahsil, forest area

Introduction

Honey bees visit plants for nectar and pollen. Nectar consisting predominantly of sources often associated with limited quantity of glucose and pollen grains provide the chief source of protein requirement of the bees essential for building their body tissues. (Rahman Khan 1941) [10] particularly during the early embryonic growth, bees prefer the nectar of a plant species that has the maximum sugar concentration. (Ramanujam 1991) [11] Similarly they prefer pollen type with the maximum nutritive values and palatability. Melittopalynological investigation involving honey samples and pollen loads furnish reliable information on the relative preferences of the honey bees among the floral sources available within their foraging ranges. (Ramanujam 1994) [12] Analysis of pollen load unravels the floral fidelity of fixity of the bees to a particular plant species in any floristic community, by highlighting the numerical status of the pollen type in the individual loads. The quantification of the data would help us to recognize the major and minor sources of pollen in any particular area. (Chaudhari 1978) [3]

Studies involving the analysis of pollen loads are few when compared to those of honeys, in the Indian context. Sharma

(1970 a & 1970 b, 1972) [6, 7, 8] and Chaturvedi (1973) studied the pollen loads of *Apis cerena*, the Indian hive bee, from Kangra in Himachal Pradesh and Banthara in the vicinity of Lucknow. Seethalakshmi and Perey (1980) [9] recognized *Borassus flabellifer* as a good pollen source in Tamilnadu by analysing 900 pollen loads of *Apis cerena* at Vijayarai in West Godavari District of Andhra Pradesh and recognized potential of this region for apiculture Kalpana, Khatija and Ramanujam (1990) [13] and Ramanujam and Kalpana (1990) [14] provided information on the pollen sources of *Apis florea* and *Apis cerena* honey bees in Hyderabad and Ranga Reddy District. Recently Borkar Laxmikant and Mate Devendra (2014) [1] provided information on the pollen source of *Apis dorsata* Honeybees in the Bramhapuri forest area of Chandrapur District of Maharashtra state and Cherian *et al.* (2011) [2] provided information on the pollen sources of *Apis cerena* honeybees in Nagpur District of Maharashtra. This study is aimed to recognize the major and minor sources of pollen to *Apis dorsata* bee in these forest during summer period (Honey flow season) on the basis of qualitative and quantitative analysis of numerous pollen loads recovered directly from various honeycombs.




Fig 1: Map Showing Chandrapur Tahsil of Chandrapur district from where the pollen loads were collected

Material and method

Pollen loads (Comb loads) 76 in number of *Apis dorsata* were obtained from two Honeycombs collected on 16 March 2012 to 3 May 2013 from Lohara and Chichpali forest area of Chandrapur tahsil of Chandrapur District of Maharashtra State. (CHN-CHN-LOH), (CHN-CHN-CHI).

The pollen grains of each pollen load were dispersed in 1 ml of glacial acetic Acid and later on subjected to acetolysis. Erdtman (1960) [4] One slide prepared for each pollen load and microscopically examined. All such pollen loads consisting of a single pollen type represent unifloral loads, with two pollen types bifloral and with more than two, multifloral Sharma, (1970 a) [6]. Identification of the pollen types was based upon the reference palynoslides of the forest flora and the relevant literature. The pollen

productivity of the significant taxa was computed using haemocytometer.

Result

The analysis has brought to light that 8 (10.52 %) loads were unifloral, 23 (30.26 %) were bifloral and the remaining 45 (59.21 %) loads multifloral (Table 2).

The pollen grain of 12 taxa referable to 9 families were recorded. These are *Terminalia* sp. (Combrataceae), *Mangifera indica* (Anacardiaceae), *Blumea* sp. (Asteraceae), *Delonix regia* (caesalpinaceae), *Bombax ceiba* (Bombiaceae), Of these *Blumea* sp. And *Carthamus tinctorius* are herbaceous weeds which represent the undergrowth, the remaining taxa are either arborescent member or shrub of the forest range.

Table 1: Pollen morphological characters of the Taxa recorded

S.N.	Pollen Type	Size, Shape & Symmetry	Aperture Pattern	Pollen Wall (sporoderm) structure & sculpture
Combrataceae				
01	<i>Terminalia</i> sp	19-22 µm, Amb spheroidal; 21-24 x 20-22 µm, subprolate; Radially symmetrical	Tricolporate, colpi alternating with pseudocolpi colpi linear, tips acute pseudocolpi almost equal the size of colpi, ora more or less circular	Exine 1.5 µm thick, tectae, surface psilate to locally finely granular
Anacardiaceae				
02	<i>Mangifera indica</i> Linn.	27-31 µm, Amb subtriangular; 29-32 x 26-28 µm, subprolate; Radially symmetrical	Tricolporate colpi long, tips acute ora prominently lanlongate	Exine 2.5 µm thick, subtectate, surface striatoreticulae, striations more or less parallel in equatorial view, lumen generally elongated in polar direction, murisimplibaculate
Asteraceae				
03	<i>Tridax procumbens</i> Linn.	31-38 µm, Amb rounded triangular to squarish; 30-35 x 32-38 µm, oblate spheroidal; Radially symmetrical	Tri to tetra colporate, colpi linear, sharply tapering, ora faint, circular	Exine 5 µm (without spines) thick, tectate, surface echinate, spines 6 µm long, 2.5 µm in diam, at base
04	<i>Blumea</i> sp.	21-24 µm, Amb spheroidal, isopolar, Radially symmetrical	21-24 µm, Amb spheroidal, isopolar, Radially	Exine 3 µm thick, surface echinate, spines 5-6 µm long, 4 spines in the interapertural region interspinal area psilate
05	<i>Lagascea mollis</i> Cav.	38-42 µm, Amb spheroidal to rounded triangular; 33-35 x 39-43 µm, oblate spheroidal; Radially symmetrical	Tricolporate, colpi linear, tips acute, ora lalongate	Exine 5 µm thick tectate, surface echinate, spines 6.5 µm long, base 2.3 µm broad
06	<i>Carthamus tinctorius</i> Linn.	59-65 µm, Amb spheroidal: 58-62 x 66-73 µm, subprolate, radially symmetrical	Tricolporate, colpi with tapering ends, ora lalongate	Exine (spinoid processes included) about 8 µm thick at poles, 10 µm at equator tectate, tectum prominently columellate, columella simple or branched, sharply undulating with supractectal solid, pointed, robust sinule like processess
Meliaceae				
07	<i>Azadirachta indica</i> A.juss	50-54µm, Amb squarish, sides convex; 47-54 38-47µm, subprolate, poles smoothly rounded; Radially symmetrical	Tetracolporate, colpi long, ends tapering, tips acute, ora lalongate	Exine 3 µm thick, tectate, surface psilate to locally granular
Mimosaceae				
08	<i>Prosopis juliflora</i> (Sw.) DC	36-39 µm, Amb rounded triangular; 38-42 x 30-35 µm, prolate to subprolate; Radially symmetrical	Tricolporate, occasionally syncolpate, colpi tapering towards poles, tips acute, ora lalongate	Exine 3.2 µm thick, tectate surface faintly reticulate
Lecythidaceae				
09	<i>Careya arborea</i> Roxb.	52.1 x 40.1 µm (48-54 x 37.5-43.5) µm, subprolate, isopolar, radially symmetrical	Hexacolpate, syncolpate with crassimarginate colpi, col. Length 43.5 (42-46.5) µm	Exine thick, 3 µm, undulating, considerable thick at the poles sexine-nexine not differentiated medium reticulate, more coarse at the poles.
Bombiaceae				
10	<i>Bombax ceiba</i> Linn	51 µm (49.5 x 52.5) µm,	Tricolpate, col. length 12 (10.5-13.5)	Exine thick 3 µm, coarsely reticulate, mesh

		peroblate, isopolar, Radially symmetrical	µm	4.1 µm (3-4.5 µm) in the major part except at the angles showing medium reticulations 1-8 µm (1.5 -3 µm), greater number of baculae are found in the lumen. Muri simplibaculate, faint LO pattern.
Rutaceae				
11	<i>Citrus Sp.</i>	27-29 µm, Amb squarish, 26-30 ×25-27 µm, prolate spheroidal radially symmetrical	Tetracolporate, colpi linear, tips acute, ora lalongate	Exine 2 µm thick subtectate, surface Reticulate. Heterobrochate, meshes smaller near the apertural regions and larger elsewhere, lumina hexa to pentagonal or irregular, psilate, muri simpli to locally duplibaculate
Fabaceae				
12	<i>Pongamia pinnata</i>	29-31 µm, Amb subtriangular: 27-31× 25-28 µm, subprolate; Radially symmetrical	Tricolporate, colpi linear to narrowly elliptic tips acute, ora lalongate	Exine 1.5 µm thick, subtectate, surface granular to locally faintly microreticulate

The unifloral pollen loads include 8 unifloral loads 4 each of (10.52%) of *Terminalia sp.*, 4 (12.5%) of *Mangifera indica* (Fig.1) and bifloral 23(30.26%) include *Terminalia sp.*, *Mangifera indica*, *Blumea sp.*, *Prosopis juliflora*, *Delonix regia*, *Careya arboreya*, *Azadirachta indica*, in combination.

The multifloral loads which are encountered showed the pollen types of *Terminalia sp.*, *Mangifera indica*, *Azadirachta indica*, *Blumea Sp.*, *Delonix regia*, *Careya*

arboreya, *Bombax ceba*, *Prosopis juliflora*, *Tridax procumbens* and *Citrus sp.* (Fig. 2).

When the representation (Irrespective of percentage) of the various pollen types in the total number of pollen loads studied was considered & the percentages of pollen types recorded in each bifloral and multifloral loads were determined by counting 200 pollen grains at random, (Sharma 1970a) [6] pollen of *Terminalia sp.* were noted in as many 66 loads (86.84%) followed by *Mangifera indica* in 65 loads (85.52%).

Table 2: Analysis of pollen loads from honeycomb

Chandrapur Tahsil							
Comb	Total Po Pollen Loads	Unifloral Loads		Bifloral Loads		Multifloral Loads	
		Number	Composition	Number	Composition	Number	Composition
CHN-CHA-Chi-25	42	03	2 – Ma 1 – Te	10	7-Ma(12,65), Te(35,88) 2-Te(65,66), Bl(34,35) 1-Ma(72), Az(28)	29	8-De(8,61), Te(16,76), Ma(8,10), Bl(6,15) 7-Te(67,85), Ma(7,20), Bl(8,13) 5-Car(3,25), Bo(10,14), Te(50,57), Ma(11,30) 3-Te(37,41), Ma(27,30), Bl(19,20), Tri(10,16), 2-Te(45,86), Ma(8,51), Ci(4,6), 1-Te(49), Ma(29), Bl(22), 1- Ma(79), De(15), Pon(4), Ps(2), 1-Te(74), Bo(12), Cart(14) 1-Te(87), Pr(9), La(4)
CHN-CHA-Loh-11	34	05	3 – Te 2 – Ma	13	12-Te(15,79), Ma(21,85) 1-Ma(61), Pr(39)	16	6-De(6,8), Ma(22,32), Te(60,72) 5-Bl(9,11), Ma(21,39), Te(50,70) 2-Pr(12,50), De(45,76), Ma(2,15) 2-Te(53,77), Bo(3,45), Bl(2,20) 1-Bo(11), Pr(27), Te(58), Ci(4)
Total	76	8 (10.52%)		23 (30.26%)		45 (59.21%)	

Abbreviations for pollen types recorded from pollen loads

Te- *Terminalia sp.* Ma- *Mangifera indica* Bl- *Blumea sp.* Ci- *Citrus sp.* Cart- *Carthamus tinctorius* Bo- *Bombax ceiba* Pr- *Prosopis juliflora* Tri- *Tridax procumbens* Pon- *Pongamia pinnata* Az- *Azadirachta indica* Car- *Careya arborea* La- *Lagascea mollis*



Fig 1: Pollen types in unifloral Pollen Loads




Fig 2: Light Microscopic photograph of pollen grain in pollen loads

Discussion

The analysis showed that the pollen loads obtained from the beehives of *Apis dorsata* in the Lohara and chichpali forest area of Chandrapur Tahsil of Chandrapur District of Maharashtra State, originated predominantly from some of the characteristics arborescent and shrubby plants of this forest area. Viz. *Terminalia* sp, *Mangifera indica*, *Delonix regia*, *Prosopis juliflora*, *Bombax ceiba*, *Blumea* sp. The

contribution to herbaceous weeds such as *Blumea* sp., *Tridax procumbens* as pollen source to *Apis dorsata* bees is very meagre.

The quantification of the data reveals unequivocally the predominance of the pollen of *Terminalia* sp as evidenced by its very high representation of 50 % in the Unifloral loads and 86.84% in the totality of the pollen loads material studied.

It can therefore be concluded that *Terminalia* sp constitutes the major source of pollen to the honey bees during the summer period. The other fairly significant source of pollen to the honeybees of this area are *Mangifera indica* 65 (85.52%), *Delonix regia* 17 (22.36 %), *Bombax ceiba* 9 (11.84%).

All these taxa also constitute important pollen source during the summer season for the honeybees of this forest area.

References

1. Borkar Laxmikant, Mate Devendra. Summer Pollen Sources to *Apis dorsata* honey bees collected from Bramhapuri forest area of Chandrapur District of Maharashtra State (India). Int. j. of life sciences, 2014;2(2):160-164.
2. Cherian KJ, Bhowal M, Godghate SD. pollen and physico-chemical analysis of honey produce by *Apis Cerena indica*. of Nagpur, Maharashtra india. Journal of Environmental Research and Developmen, 2011;5(3): 542-550.
3. Chaudhari RK. Floral fidelity in the Indian honey bee (*Apis cerana indica* f.) Indian Bee journal, 1978;49(2):33-35.
4. Erdtman G. The acetolysis method a rivised descreption seven -Botan.Tidskr., 1960:54:561-564.
5. Kalpna TP, Khatija f, Ramanujan CGK. pollen anlaysis of *Apis Cerena* and *Apis florea*. honeys, from Adikmet. area, Hyderabad prc.ind. Acad Sci. (plant Sci), 1990:100 (3):183-193.
6. Sharma M. An analysis of pollen loads of honeybees from kangra, india *Grana*, 1970a:10:35-42.
7. Sharma M. Studies on the pollen loads of honeybees from kangra india Journal of palynology, 1970b:6:104-110.
8. Sharma M. Studies on the pollen loads of honeybees from kanga india, tour Poly, 1972:66:04:110.
9. Seethalakshmi TS, Percy AP. Borassus flabellifer (palmryrah Pollen) a good pollen source. Indian bee Journal, 1980:41(1-2):20-21.
10. Rahman Khan A. Nectar and pollen plants of the Punjab, Indian Bee journal, 1941:4:32-35.
11. Ramanujam CGK. Beekeeping & melitopalynology perspectives & prospectus.pro. T.Navaneeth Rao Commemoration volume, osmania, University, Hyderabad, 1991, 74-79.
12. Ramanujam CGK. forage sources for Rock bees during May to July in deciduous forest of Ranga Reddy District, A.P. *Gephytology*, 1994:24(1):119-122.
13. Ramanujam CGK, Kalpana TP. Pollen analysis of *prosopis juliflora* honey from Ranga Reddy District, A.P. and its relevance to agriculture and social forestry tour paly, 1990:26-28:345-368.
14. Ramanujan CGK, Fatima Khatija. summer pollen sources to "*Apis dorsata*" honeybee in deciduous forest of mahaboobnager district Andra Pradesh, *Geophytology*, 1992:21:155-161.