



Effect of integrated nutrient management on grain yield, straw yield and harvest index of transplanted rice

Biswanath Ganguly, V Imayavaramban, G Murugan

Department of Agronomy, Faculty of Agriculture, Annamalai University, Annamalai Nagar, Tamil Nadu, India

Abstract

The field experiments were conducted to evaluate the effect of inorganic fertilizers along with organic manures at the Experimental Farm, Department of Agronomy, Annamalai University, Tamil Nadu. The treatments were tested in Split-plot Design and replicated thrice. It was observed that the grain yield, straw yield and harvest index of rice crop influenced by integrated application of chemical fertilizers and organic manures. Application of 150% of recommended dose of nitrogen, phosphorus and potassium + 75% of recommended dose of Vermicompost @ 3.75 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT recorded higher grain, straw yield and harvest index. This was on par with 125% of recommended NPK + 75% of recommended dose of Vermicompost @ 3.75 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT.

Keywords: grain yield, straw yield, harvest index, INM.

Introduction

Rice, one of the predominant member of the graminea family. It is the staple food for more than 50 per cent of world population. Globally, it plays a vital role in food as well as nutritional security for several billions of human beings and thus the slogan "Rice is Life" coined by IRRI during 2004 seems to be the most appropriate (Chandrasekaran *et al.*, 2007) ^[10]. In India, rice cultivated in an area of 44.0 million ha⁻¹ with a production of 116.8 million tonnes and the average productivity of 2.93 t ha⁻¹. However, considering the current population growth rate of 1.5 per cent the projected demand for rice for the exploding, population the total rice requirement by 2025 would be around 300 million tonnes (Panda, 2010) ^[16]. Although sixteen numbers of nutrients constituted of macro, secondary and micronutrients are required for optimal growth and establishment of crop plants, the major nutrient elements *viz.*, nitrogen, phosphorus and potassium are essentially needed for varied metabolic process of crop plants, while nitrogen is the key nutrition synthesis of ATP. It is the prime component of chlorophyll, proteins and enzymes and it also helps the plants in synthesis and utilization of CHO (Mengel and Kirkby, 2001; Sara *et al.*, 2013) ^[14, 18]. P is a part of nucleotides like present in Adenosine diphosphate and Adenosine triphosphate energy bonds, it involved in many processes such as mitotic activities, tissue growth and development (Bhattacharyya and Jain, 2000) ^[5]. The crop plants could not achieve its maximum yield potentials without supply of adequate phosphorus (Alam *et al.*, 2009; Murtaza *et al.*, 2014) ^[2, 15]. It also play a crucial role in root proliferation, consistent grain filling and quality. Similarly, K is vital for the supply of electrical potential across cellular membranes and cellular turgor, which permits cell expansion and enlargement, opening and closing of stomata and pollen tube development. Further, it is also an activator of many enzymes and it also involved in translocation of nitrate and sucrose (Britto and Kronzucker, 2008) ^[9]. The farmyard manure (FYM) is the only available bulky organic manure which is being utilized for crop production by the farmers for the past several decades. It is a mixture of cattle dung, urine, litter or bedding material, portion of fodder not consumed by cattle and domestic wastes. Application of farmyard manure improves the soil fertility (Gora *et al.*, 2017) ^[11]. Its incorporation, significantly improves soil physico-chemical characteristics, prevents soil erosion improves soil organic carbon content, enhances microbial activity, neutralizers soil pH, minimizes the electrical conductivity and furthermore the decomposed FYM, act as an 'chelate', which would retains the nutrient elements within its biomass and release them as and when the crop require. However, is becoming scarcely available, due to drastically dwindling cattle population. However, its timely inavailability escorted cost factor, the cumbersome and the extended time involvement for its transportation limits its usage. Compostvermi is the decomposed product of bio solid wastes by earthworms gut and secreted as casts (Janagan *et al.*, 2003) ^[13]. It contains most of the macro and micronutrients that too in easily available forms facilitating enhanced crop uptake growth and productivity as well as maintaining up of soil health (Marinari *et al.*, 2008). Pressmud is an one of the most valuable organic by-product obtaining from the sugar industry, before crystallization of sugar from sugar juice (Agarwal *et al.*, 2014).

Biofertilizers are eco-friendly, cost effective organic agro inputs which are essential nutritional supplement for crop plants in conjunction with wide range of beneficial enzymes and growth hormone (Bloemberg *et al.*, 2000; Amudha *et al.*, 2014) ^[7, 3]. From ancient times, seaweeds is also considered to be source of food, fodder, medicine and an nutrients supplementing bio-inoculant. The commercial product of seaweed liquid fertilizer (SLF) is available which would enhance the productivity of crop plants as an supplemental nutrition supply (Bokil *et al.*, 1974 and Bai *et al.*, 2007) ^[8, 4].

Materials and Methods

The field experiment was conducted to study the effect of INM on rice at Experimental Farm, Department of Agronomy, Annamalai University with rice var. CO-51. The research comprised of

A. Main: Plot (NPK doses)

M₁: 100% of recommended NPK

M₂: 125% of recommended NPK

M₃: 150% of recommended NPK

B. Sub: Plot (Organic inputs levels)

S₁: 100% of recommended dose of FYM @ 12.5 t ha⁻¹

S₂: 75% of recommended dose of FYM @ 9.5 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla @ 10 kg ha⁻¹ on 10 DAT + foliar application of seaweed extract @ 2.0% on 15 and 30 DAT.

S₃: 50% of recommended dose of FYM @ 6.25 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla @ 10 kg ha⁻¹ on 10 DAT + foliar application of seaweed extract @ 2.0% on 15 and 30 DAT.

S₄: 75% of recommended dose of Vermicompost @ 3.75 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla @ 10 kg ha⁻¹ on 10 DAT + foliar application of seaweed extract @ 2.0% on 15 and 30 DAT.

S₅: 50% of recommended dose of Vermicompost @ 2.50 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla @ 10 kg ha⁻¹ on 10 DAT + foliar application of seaweed extract @ 2.0% on 15 and 30 DAT.

S₆: 75% of recommended dose of fortified pressmud @ 3.75 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla @ 10 kg ha⁻¹ on 10 DAT + foliar application of seaweed extract @ 2.0% on 15 and 30 DAT.

S₇: 50% of recommended dose of fortified pressmud @ 2.5 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT.

Results and Discussion

Grain yield (Table 1)

Grain yield of rice was significantly influenced by the integration of nutrients through chemical fertilizers and organic manures in the experiment. Among the different levels of inorganic fertilizers, application of 150% of recommended NPK (M₃) recorded highest grain yield of 6688 kg ha⁻¹ in experiment. Application of 100% of recommended NPK (M₁) registered lower grain yield of 5958 kg ha⁻¹ in experiment. The effects between the inorganic fertilizer levels and organic manure sources, application of 75% of recommended dose of Vermicompost @ 3.75 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (S₄) appear its dominance by producing the highest grain yield of 6901 kg ha⁻¹ in experiment. This was statistically on par with the application 75% of recommended dose of FYM @ 9.5 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (S₂). The lowest grain yield of 6617 kg ha⁻¹ in experiment was recorded with the application of 50% of recommended dose of fortified pressmud @ 2.5 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (S₇). Similarly, application of 150% of recommended NPK + 75% of recommended dose of Vermicompost @ 3.75 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla @ 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (M₃S₄) excelled all other treatments and registered the highest grain yield of 7252 kg ha⁻¹ in experiment, respectively. The lowest grain yield of 5578 kg ha⁻¹ in experiment was observed in the combined application of 100% of recommended NPK + 50% of recommended dose of fortified pressmud @ 2.5 t ha⁻¹ + Azophos at the rate of 10 kg ha⁻¹ + *in situ* application of Azolla @ 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (M₁S₇).

Table 1: Effect of NPK doses and the levels of manures and other inputs on grain yield (kg ha⁻¹)

Treatments	Grain yield (kg ha ⁻¹)			
	M ₁	M ₂	M ₃	Mean
S ₁	6053	6564	6788	6468
S ₂	6242	6655	6956	6617
S ₃	5555	6057	6427	6013
S ₄	6556	6895	7252	6901
S ₅	5765	6292	6477	6178
S ₆	5962	6459	6690	6370

S ₇	5578	6034	6226	5946
Mean	5958	6422	6688	–
SEd			CD (P=0.05)	
M	287.4		604.7	
S	251.1		527.3	
M × S	214.8		452.0	
S at M	215.6		453.8	

Straw Yield (Table 2)

Integrated nutrient management practices exerted marked influenced on straw yield of rice at harvest stage. Among the different levels of inorganic fertilizers, application of 150% NPK (M₃) recorded highest straw yield of 7807 in experiment at harvesting stage. Application of 100% of recommended NPK (M₁) registered lowest straw yield of 7400 kg ha⁻¹ experiment at harvesting stage. Significant impact on straw yield of growth were documented with the application of different levels of manures and other organic inputs in both the years of experimentation. Application of 75% of recommended dose of vermicompost @ 3.75 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (S₄) significantly registered highest straw yield of 7678 kg ha⁻¹ experiment at harvest stage. The lowest straw yield of 7565 kg ha⁻¹ was resulted with application of 50% of recommended dose of fortified pressmud @ 2.5 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (S₇) in experiment at harvest stage. Similarly, application of 150% of recommended NPK + 75% of recommended dose of Vermicompost @ 3.75 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (M₃S₄) excelled all other treatments and registered the highest grain yield of 7896 kg ha⁻¹ in experiment. The lowest grain yield of 7385 kg ha⁻¹ in experiment were recorded in the combined application of 100% of recommended NPK + 50% of recommended dose of fortified pressmud @ 2.5 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (M₁S₇).

Table 2: Effect of NPK doses and the levels of manures and other organic inputs on straw yield (kg ha⁻¹)

Treatments	Straw yield (kg ha ⁻¹)			
	M ₁	M ₂	M ₃	Mean
S ₁	7402	7406	7920	7576
S ₂	7420	7653	7987	7686
S ₃	7204	7370	7615	7396
S ₄	7530	7609	7896	7678
S ₅	7466	7501	7808	7591
S ₆	7398	7608	7604	7536
S ₇	7385	7489	7823	7565
Mean	7400	7519	7807	–
SEd			CD (P=0.05)	
M	95.6		201.3	
S	70.7		148.7	
M × S	86.6		182.2	
S at M	86.9		183.0	

Harvest Index (Table 3)

Integrated nutrient management practices exerted marked influenced on harvest index of rice at harvest stage. Among the different levels of inorganic fertilizers, application of 150% of recommended NPK (M₃) recorded highest harvest index of 46.1 in experiment at harvesting stage, respectively. Application of 100% of recommended NPK (M₁) registered lowest harvest index of in 44.5 experiment at harvesting stage. Similarly, application of 150% of recommended NPK + 75% of recommended dose of Vermicompost @ 3.75 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (M₃S₄) excelled all other treatments and registered the highest harvest index of 47.3 in experiment. The lowest harvest index of 46.2 in experiment were recorded in the combined application of 100% of recommended NPK + 50% of recommended dose of fortified pressmud @ 2.5 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (M₁S₇). Among the main plot treatments, application of 150% of recommended NPK (M₃) showed a significant influence on grain and straw yield. This might be due to application of 150% of recommended NPK in the crop and it has enhanced crop grain and straw yield production. This result is in conformity with the reports of (Sarker *et al.* 2008) ^[19]. With respect to subplot treatments, a positive response of growth parameters was noticed to different organic manures combination of nutrients, irrespective of the different combination of nutrients. The response was observed upto 75% of

recommended dose of Vermicompost @ 3.75 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (S₄) has significantly increased grain and straw yield. The integrated application of chemical and organic manures increased the grain yield and straw yield of rice led to increase the growth and yield attributes. The effective application of nutrient delayed the senescence and increased the life span of the plant and it resulted in higher economic yield. The sustainable use of nuclear and chemical fertilizer enabled the rice plant to assimilate sufficient photosynthates resulting in increased dry matter production and these together produced more productive tillers, panicle and number of filled grains leading to higher grain yield, these results in accordance with (Hossaen *et al.*, 2011) [12]. Among the various treatment combinations, 100% of recommended NPK + 75% of recommended dose of Vermicompost @ 3.75 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (M₃S₄). Which efficiently increased the grain yield when compared to other treatment combinations. The above treatment also recorded higher straw yield due to higher LAI and DMP. These results are in conformity with the findings of Singh *et al.* (2005) [21]. Within in the plant treatments, application of 150% of recommended NPK (M₃) showed a significant ascendancy on harvest index. It is due to the application of 150% of recommended NPK in the crop plant and it enhanced into the crop yield attributes. This results are ascertained with the (Rao *et al.*, 2020) [17]. With respect to sub plot treatments incorporation of 75% of recommended dose of vermicompost @ 3.75 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla @ 10 kg ha⁻¹ on 10 DAT + foliar application of seaweed extract @ 2.0% on 15 and 30 DAT (S₄) significantly registered higher harvest index of 47.3%. The lowest harvest index of 43.9% experiment at harvest stage was resulted with the incorporation of 50% of recommended dose of fortified pressmud @ 2.5 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (S₇) treatment (Biramo, 2018) [6]. The increased in harvest index percentage depends on the developmental stages of the crop (grain and straw yield). Application of integrated nutrient management in a efficient amount leads towards the higher yield and it proportionally increased harvest index percentage with reflect on economics as well. These results are in conformity with the findings of Singh *et al.* (2018) [22]. Among the various treatment combinations, 150% of recommended NPK + 75% of recommended dose of Vermicompost @ 3.75 t ha⁻¹ + Azophos @ 10 kg ha⁻¹ + *in situ* application of Azolla at the rate of 10 kg ha⁻¹ on 10 DAT + foliar nutrition of seaweed extract @ 2.0% on 15 and 30 DAT (M₃S₄). Which efficiently increased the harvest index percentage when compared to other treatment combinations. The above treatment also recorded higher harvest index percentage due to higher grain and straw yield. These results are in similar with the reports of Tomar *et al.* (2015).

Table 3: Effect of NPK doses and the levels of manures and other organic inputs on harvest index (%)

Treatments	Harvest index (%)			
	M ₁	M ₂	M ₃	Mean
S ₁	44.9	46.9	46.2	45.9
S ₂	45.6	46.5	46.6	46.2
S ₃	43.5	45.1	45.7	44.7
S ₄	46.5	47.5	47.9	47.3
S ₅	43.2	45.6	45.3	44.8
S ₆	44.7	45.9	46.8	45.8
S ₇	43.1	44.6	44.3	43.9
Mean	44.5	46.0	46.1	—
SEd			CD (P=0.05)	
M	0.59		1.24	
S	0.69		1.45	
M × S	0.87		1.85	
S at M	0.88		1.87	

Conclusion

On the basis of experiment finding the conclusions that INM proved its superiority in increasing the grain yield, straw yield and harvest index percentage under sustainable crop productivity.

References

1. Agrawal KM, Barve BR, Khan SS. Biogas from pressmud. IOSR J. Mech. Civil Eng,2014;2:37-41.
2. Alam MM, Ali MH, Amin AKMR, Hasanuzzaman M. Yield attributes, yield and harvest index of three irrigated rice varieties under different levels of phosphorus. Adv. Biol. Res,2009;3:132-139.
3. Amudha RS. Karunakaran S, Dhanasekaran K. Hemalatha R, Monika Shanmugapriya T. Pand Sornolatha. Isolation and mass production of biofertilizers (Azotobactor and Phosphobacter). Int. J. Lat. Res. Technol,2014;3:79-81.
4. Bai NR, Banu NRL, Prakash JW, Goldi SJ. Effects of Asparagopsis taxiformis extract on the growth and yield of (Phaseolus aureus). J. Basic Appl. Biol,2007;1:6-11.

5. Bhattacharyya P, Jain RK. Phosphorus solubilizing biofertilizers in the whirlpool of rock phosphate-challenges and opportunities. *Fert. News*,2000:45:45-51.
6. Biramo G. The role integrated nutrient management system for improving crop yield and enhancing soil fertility under small holder farmers in sub-Saharan Africa : A Review Article. *J. Nat. Sci. Res*,2018:8:26-35.
7. Bloemberg GV, Wijfij AHM, Lamers GEM, Stuurman N, Lugtenberg BJJ. Simultaneous imaging of *Pseudomonas fluorescence* WCS 3655 populations expressing three different auto fluorescent proteins in rhizosphere: new perspective for studying microbial communities. *Mol. Plant Mic. Int*,2000:13:1170-1176.
8. Bokil KK, Mehta VC, Datar DS. Seaweeds as manure: II pot culture manorial experiments on wheat. *Phykos*,1974:13(1):1-5.
9. Britto DT, Kronzucker HJ. Cellular mechanisms of potassium transport in plants. *Physiol. Plant*,2008:133:637-650.
10. Chandrasekaran B, Annadurai K, Kavimani R. A textbook of rice science. Scientific publishers (India), Jodhpur, India,2007, 7.
11. Gora MK, Choudhary RR, Sharma OP, Shankar Lal Choudhary. Effect of organic manures and molybdenum on growth, yield and quality of Groundnut. *Int. J. Curr. Microbiol. Appl. Sci*,2017:6(6):736-742.
12. Hossaen MA, Shamsuddoha AT, Paul AK, Bhuiyan MSI, Zobaer ASM. Efficiency of different organic manures and inorganic fertilizer on yield and yield attributes of boro rice. *The Agric*,2011:9(1&2):117-125.
13. Janagan KV. Sathish, Vijaykumar A. A sustainable system for solid waste treatment-Vermiculture Didalam: Bunch, M.J., Suresh, V.M., Kumaran, T.V. (ed). *Proceedings of the third international Conference on Environment and Health*, Chennai, India, 15-17 December, 2003. Chennai: Department of Geography, University of Madras and Faculty of Environmental studies, York University, 2003, 175-185.
14. Mengel K, Kirkby EA. Boron: In principles of plant nutrition, 5th ed. Kluwer Academic Publishers (5th ed.) Dordrecht, Boston, London, Netherlands, 2001.
15. Murtaza G, Zohaib A, Hussain S, Rasool T, Shehzad H, Ehsanullah. The influence of rhizobium seed inoculation and different levels of phosphorus application on growth, yield and quality of mashbean (*Vigna mungo* L.). *Intl. J. Mod. Agri*,2014:3:92-96.
16. Panda BB. Biochemical studies on iron metabolism in rice plant (*Oryza sativa* L.). Ph.D. Thesis, Sambalpur University, Sambalpur, 2010.
17. Rao A, Singh NB, Pandey D, Singh MP. Effect of integrated nutrient management on productivity and profitability of rice under SRI. *J. Pharmacog. Phytochem*,2020:9:665-667.
18. Sara S, Morad M, Reza CM. Effects of seed inoculation by Rhizobium strains on chlorophyll content and protein percentage in common bean cultivars (*Phaseolus vulgaris* L.). *Intl. J. Biosci*,2013:3:1-8.
19. Sarker ABS, Islam MA, Islam MR. Effect of phosphorus on nutrient uptake of *Japonica* and *Indica* rice. *J. Agric. Rural. Dev*,2008:6(1&2):7-12.
20. Singh NP, Singh MK. Tyagi S, Singh SS. Effect of integrated nutrient management on growth and yield of rice (*Oryza sativa* L.). *Int. J. Curr. Microbiol. App. Sci*,2018:7:3671-3681.
21. Singh Y, Sharma HC, Singh K, Singh UN, Singh CS, Singh KK. Rice yield as affected by use of bio fortified straw compost combined with NPK fertilizers. *Int. Rice Res. News*,2005:30:47-48.
22. Tomar R, Singh NB, Singh V, Kumar D. Effect of planting methods and integrated nutrient management on growth parameters, yield and economic of rice. *J. Pharmacog. Phytochem*,2018:7:520-527.