

Hydrobiological studies on Narsarha Talab of Shahdol Madhya Pradesh

Tasleem Begum

Guest Faculty, Dept. of Botany, Govt. P.G. College, Bareilly District Raisen, Madhya Pradesh, India

Abstract

The water quality has become a major concern due to over increasing human development activities that over exploits and pollute the water resources. Present investigations were carried out on the limnological aspects of Narsarha Talab in Shahdol district (M.P.). Many of the parameters were found below the permissible limits for drinking water as suggested by WHO. A total of 11 parameters were analyzed and their seasonal variations in the year 2014-2015 were discussed.

Keywords: Physico-chemical, Limnology, Narsarha Talab, Shahdol district, Seasonal variations

1. Introduction

The aquatic ecosystem is extremely important to mankind as they have various uses, including drinking water supply, irrigation, navigation, recreation etc. and are also source of organic productivity. During recent year there has been increasingly great concern for inland freshwater resources which are affected in different ways by all kinds of anthropogenic activities. Therefore the limnological investigations on water bodies are need of today. Several studies have been made on the limnology of fresh water bodies in India (Ganapathi, 1940; Alikunhi *et al.*, 1948; Harshey, *et al.*, 1982; Rao and Mahmood, 1995; Naganandini and Hosmani, 1998; Patel and Sinha, 1998; Kashyap, 2015) [4, 1, 6, 15, 13, 14, 8]. But unfortunately very little information is available on the limnology of this region. Therefore, the present study was undertaken of Narsarha Talab in Shahdol district (M.P.).

1.1 Study Area

The Shahdol district lies in the north-east part of Madhya Pradesh extending from 29°39'28" and 24°16'13" North latitude and from 80°32'56" to 82°12'21" East longitude approximately. The region lies in the heart of the country. The district is surrounded by Sone river and Rewa district in North, Mandla, in South Sidhi, Anuppur in Surguja East and Umaria and Satna in West. It is situated 489 meter above of the sea surface.

The Shahdol district has undulating and hilly topography. The southern-most part of the district is hilly and form part of the Maikal range, while the portion north of it is quite undulating. In the south most portion has hills of higher elevation as compared to northern portion. There is a continuous drop in elevation from south to north and it attains its lower most elevation in the extreme north where the some forms the boundary of the district. The southernmost portion (parts of south and north Shahdol divisions) forming part of Maikal range is generally more than 900 m. above msl, while the portion north of it seldom reaches 800 m. msl.

The present aquatic body is perennial one. The area of talab is nearly 10 hectare. It is situated within municipal area of Shahdol city. The fish culturing carried out since last ten years. Hence it has been selected for the present study.

2. Materials and Methods

To study the water quality and its seasonal variations, the water samples are collected from the surface at a depth of 22 cm. from four different points, integrated and a representative sample was taken. Water samples were collected during morning hours in between 8.30 to 10.30 a.m. with one liter containers from the dam in three seasons i.e. during summer, monsoon and winter seasons from April 2014 to March 2015. Sampling was done at four sites. Some of the results were recorded at the sampling sites whereas the others were recorded in the laboratory. The parameters observed were air and water temperature, relative humidity, Dissolve Oxygen (DO), Total Dissolved Solid (TDS), Free CO₂, pH, Conductivity, and Hardness of the lake water. Samples of the water for physicochemical characteristics were analysed according to standard methods of APHA (1998) [2].

3. Result and Discussion


The results of physicochemical analysis of three seasons are summarized in Table-1. The pH of the talab is 7.22, 6.61 and 6.52 during the summer, monsoon and winter seasons. No significant change was observed in the water pH and it was found within the permissible limits of 6.5 to 8.5 (WHO). The highest value was noticed in summer season and lowest in winter season. Decline in pH during monsoon and winter seasons were attributed to the rains as it increase the amount of carbonic acid to the lake water (Khan and Chaudhary 1994, Kaushik *et al.* 1989) [10, 9]. Hutchinson (1975) [7] stated that if any aquatic system is neither higher alkaline nor highly acidic, the pH of the water is principally governed by the CO₂ - bicarbonate- carbonate system.

Table 1: Physico-chemical Parameter of Narsarha Talab in Shahdol district

| S. No. | Parameters | Seasons | | | SD |
|--------|--|------------|------------|------------|---------|
| | | Summer | Monsoon | Winter | |
| 1. | Temperature of air (°C) | 8 | 5 | 6 | ±1.522 |
| 2. | Temperature of water (°C) | 5 | 3 | 2 | ±1.527 |
| 3. | Colour | Colourless | Colourless | Colourless | - |
| 4. | pH | 7.22 | 6.52 | 6.61 | ±0.363 |
| 5. | Alkalinity mg ^l ⁻¹ | 40.00 | 36.00 | 34.00 | ±2.512 |
| 6. | TDS mg ^l ⁻¹ | 0.21 | 0.11 | 0.07 | ±0.064 |
| 7. | Hardness mg ^l ⁻¹ | 17.00 | 12.00 | 14.00 | ±1.528 |
| 8. | Chloride mg ^l ⁻¹ | 14.40 | 24.72 | 5.33 | ±10.201 |
| 9. | Total phosphate mg ^l ⁻¹ | 0.09 | 0.06 | 0.04 | ±0.023 |
| 10. | DO mg ^l ⁻¹ | 8.32 | 9.81 | 10.91 | ±1.295 |
| 11. | Free CO ₂ mg ^l ⁻¹ | 5.26 | 3.60 | 4.34 | ±0.811 |

The variation was observed in the water temperatures in three seasons whereas a well-marked seasonal variation in air temperature was recorded. Water temperature fluctuated with the air temperature, at all the sampling sites and both air and water temperature were correlated positively ($r = 0.786$) in the present study. The temperature remained comparatively low throughout the study, which may be due to the presence of thick forest around it. Martin (1972) [11]

stated that the clarity of water, presence of vegetation etc. are the factors mainly responsible for the daily fluctuations in water temperature. The alkalinity varied from 34 mg^l⁻¹ to 40 mg^l⁻¹ in three seasons, during which minimum value (34 mg^l⁻¹) was observed in winter season and the maximum (40 mg^l⁻¹) in summer season. This may be due to the liberation of CO₂ in the process of decomposition of bottom sediments with resultant conversion of carbonates to bicarbonates.

**Fig 1:** Graphics analysis of Physico-chemical Parameter of Narsarha Talab in Shahdol district

The alkalinity of water is usually caused by the presence of carbonates, bicarbonates and hydroxyl ions and less frequently by borates, silicates and phosphates (APHA 1998) [2].

Total dissolved solids of the dam were 0.21 mg^l⁻¹ in summer, which is the highest value and the lowest values 0.07 mg^l⁻¹ was noticed in winter. Total hardness value of the dam was 12 to 17 mg^l⁻¹ of which higher value was in summer while the lowest in monsoon season. The maximum permissible limit for this parameter for drinking water standards is 500 mg^l⁻¹. Chloride values were found ranging between 5.33 to 24.72 mg^l⁻¹ of which maximum value was noticed in monsoon and the lowest value in winter may be due to dilution effect in post monsoon period. Chourasia and Adoni (1985) [3] also found similar behaviour of chlorides in their studies on Sagar lake with summer maxima and winter minima. The concentration of dissolved oxygen was recorded high during winter season (10.91mg^l⁻¹). This may be due to that the oxygen concentration in the dam water

increased by the reduction in water temperature. Free Carbon dioxide is one of the essential constituents of an aquatic ecosystem. The abundance of carbon dioxide exerts certain specific effects on aquatic bioata. The dam exhibited maximum carbon dioxide as 5.26mg^l⁻¹ during summer whereas the lowest concentration of carbon dioxide 3.60mg^l⁻¹ was recorded during monsoon season. Cole (1975) noted that free CO₂ supply rarely limits the growth of phytoplankton. It is evident from the present study that the phosphate concentration was higher during summer and lower in winter season. It was quite opposite in relation to dissolved oxygen and phytoplankton population. Many earlier workers have also reported similar findings (Meckenzie and Gillespie, 1986; Ghavzan *et al.*, 2006; Kashyap, 2015) [12, 5, 8].

4. Acknowledgement

The author is thankful to authority of Dept. of Botany, Govt. P.G. College, Bareilly District Raisen (M.P.) for granting

permission to carry out this work.

5. References

1. Alikunhi KH, Ganapathi SV, Thivy F. Limnology of the Ootacamand Lake, Nilgiris II Summer conditions. Proc. 35th Ind. Sci. Congress Assoc., Part III, Calcutta, India, 1948.
2. APHA. Standard methods for the estimation of water and waste water. American public health association. American waste water association and water pollution control federation. 20th Edition Washington. DC, 1998.
3. Chourasia SK, Adoni AD. Zooplankton dynamics in a shallow eutrophic. Limnology Bot. Soc. Sagar. 1985; 32:30-39
4. Ganapathi SV. The ecology of a temple tank containing permanent bloom of Microcystic aeruginosa Kuetz. J. Bom. Nat. Hist. Soc. 1940; 42:65-77.
5. Ghavzan NJ, Gunale VR, Trivedy RK. Limnological evaluation of an urban fresh water river with special reference to phytoplankton. Pollut. Res. 2006; 25(2):259-268
6. Harshey DK, Patil SG, Singh DF. Limnological studies on a tropical fresh water fish tank of Jabalpur, India. The abiotic factors. Geobios New Reports. 1982; 1:98-102
7. Hutchinson GE. A Treatise in Limnology. John Willey and Sons Inc. New York, 1975, 1015.
8. Kashyap, Vinita R. Hydro Biological Studies on Nebuha Sam of Sidhi District (M.P.), International Education & Research Journal. 2015; 1(3):24-25. [IERJ], ISSN No : 2454-9916,
9. Kaushik S, Sharma S, Saxena MN. Aquatic insect population in relation to physico-chemical characteristics of Chandanpura pond, Gwalior (M.P.). In: Environmental Risk Management (Eds: Y.N. Sahai, P.B. Deshmukh, T.A. Mathai and K.S. Pillai) Academy of Environmental Biology, India, Muzaffernagar, 1989, 243-254.
10. Khan MAG, Chaudhary SM. Physical and chemical limnology of the lake Kaptai, Bangladesh. Tropical Ecology, 1994; 35:35-51.
11. Martin NA. Temperature fluctuations with in English lowland ponds. Hydrobiologia, 1972; 40(4):455-470.
12. Meckenzie AL, Gillespie PA. Plankton ecology and productivity nutrient chemistry and hydrography of Tasman bay. Newzeland, 1982-1984. N.Z.J. Mar. Fresh Water, Res. 1986; 20:365-395.
13. Naganandini MN, Hosmani SP. Ecology of certain inland waters of Mysore district, occurrence of cyanophycean bloom at Hosakere lake. Pollut. Res. 1998; 17(2):123 -125
14. Patel, Niroj K, Sinha BK. Study of the pollution load in the ponds of Burla area near Hirakund dam at Orissa. J. Environ. Pollut. 2000-1998; 5(2):157-160.
15. Rao VN, Mahmood SK. Nutrient status and biological characteristics of Habsiguda pond. J. Environ. Pollut. 1995; 2(i):31-34.